

4-2016

# What Moser Could Have Asked: Counting Hamilton Cycles in Tournaments

Neil J. Calkin

*Clemson University*, [calkin@clemson.edu](mailto:calkin@clemson.edu)

Beth Novick

Hayato Ushijima-Mwesigwa

Follow this and additional works at: [https://tigerprints.clemson.edu/mathematics\\_pubs](https://tigerprints.clemson.edu/mathematics_pubs)



Part of the [Mathematics Commons](#)

---

## Recommended Citation

Please use publisher's recommended citation. <http://www.maa.org/press/periodicals/american-mathematical-monthly/american-mathematical-monthly-april-2016>

This Article is brought to you for free and open access by the School of Mathematical and Statistical Sciences at TigerPrints. It has been accepted for inclusion in Publications by an authorized administrator of TigerPrints. For more information, please contact [kokeefe@clemson.edu](mailto:kokeefe@clemson.edu).

# What Moser *Could* Have Asked: Counting Hamilton Cycles in Tournaments

Neil J. Calkin, Beth Novick and Hayato Ushijima-Mwesigwa

June 3, 2015

## Abstract

Moser asked for a construction of explicit tournaments on  $n$  vertices having at least  $(\frac{n}{3e})^n$  Hamilton cycles. We show that he could have asked for rather more.

## 1 Introduction

... the cycle has taken us up through forests.

---

*Robert M. Pirsig*

In his classic book on tournaments, Moon [4, Section 10] discusses the question of exhibiting tournaments with a large number of Hamilton cycles. He poses the question (Exercise 4, attributed to Moser), of constructing a tournament on  $n$  vertices having at least  $(\frac{n}{3e})^n$  Hamilton cycles. Presumably, the intended construction is to take three tournaments,  $T_1, T_2, T_3$ , on  $\frac{n}{3}$  vertices, and construct a new tournament  $C_3(T_1, T_2, T_3)$  by orienting all edges from  $T_1$  to  $T_2$ ,  $T_2$  to  $T_3$ , and  $T_3$  to  $T_1$  (See Figure 1). The number of Hamilton cycles




Figure 1:  $C_3(T_1, T_2, T_3)$

---

*AMS Subject Classification: 05C20, 05C30, 05A16*

in  $C_3(T_1, T_2, T_3)$  which do not use any edges internal to  $T_1, T_2$ , or  $T_3$  is

$$\frac{\left(\frac{n}{3}\right)!^3}{\frac{n}{3}} \sim \sqrt{\frac{8\pi^3 n}{3}} \left(\frac{n}{3e}\right)^n > \left(\frac{n}{3e}\right)^n.$$

In this note, we show that this construction has many more Hamilton cycles. Indeed, if  $T_1, T_2$ , and  $T_3$  are all transitive, we show that the number of Hamilton cycles is asymptotic to  $\frac{1}{(1-\log 2)} \frac{(n-1)!}{(3 \log 2)^n}$ .

## 2 Background and Definitions

A *tournament* is an oriented, complete graph. A *Hamilton cycle* or *path* in a tournament  $T$ , is a spanning directed cycle or directed path in  $T$ . A tournament with no directed cycles is called *transitive*.

Counting Hamilton paths and cycles in tournaments is a very old problem, dating back to the 1940's: in one of the first applications of the probabilistic method, Szele [6] showed that the expected number of Hamilton paths in a random tournament is  $\frac{n!}{2^{n-1}}$ , therefore showing that there exists a tournament on  $n$  vertices with at least this many Hamilton paths. The same argument shows that there exists a tournament with at least  $\frac{(n-1)!}{2^n}$  Hamilton cycles. Moon observed that it seems difficult to give explicit tournaments with at least this many Hamilton cycles.

Deep results of Cuckler [3] show that every regular tournament on  $n$  vertices has at least  $\frac{n!}{(2+o(1))^n}$  Hamilton cycles.

Given tournaments  $T_1, T_2, T_3$ , we can construct a tournament  $C_3(T_1, T_2, T_3)$  by orienting all edges from  $T_1$  to  $T_2$ ,  $T_2$  to  $T_3$ , and  $T_3$  to  $T_1$ . We will call such tournaments *triangular*. Wormald [8] showed that if  $T_1, T_2, T_3$  are random tournaments, then the expected number of Hamilton cycles is  $2 \frac{(n-1)!}{2^n}$ .

We show that *all* triangular tournaments have a relatively large number of Hamilton cycles, even in the extreme case when they constructed from transitive tournaments.

Let  $S(m, k)$  denote the Stirling number of the second kind, that is,  $S(m, k)$  is the number of set partitions of  $\{1, 2, \dots, m\}$  into exactly  $k$  parts.

## 3 Main Result

**Theorem 1.** *Let  $T_1, T_2$ , and  $T_3$  be any tournaments on  $m_1, m_2$ , and  $m_3$  vertices respectively. Then the number  $H$  of Hamilton cycles in  $C_3(T_1, T_2, T_3)$  is at least*

$$H \geq \sum_{k=1}^{\min\{m_1, m_2, m_3\}} S(m_1, k) S(m_2, k) S(m_3, k) \frac{k!^3}{k}, \quad (1)$$

*with equality when  $T_1, T_2$ , and  $T_3$  are transitive.*

**Corollary 2.** If  $T_1, T_2$ , and  $T_3$  are transitive tournaments on  $\frac{n}{3}$  vertices, the number of Hamilton cycles in  $C_3(T_1, T_2, T_3)$  is asymptotic to

$$\frac{1}{(1 - \log 2)} \frac{(n-1)!}{(3 \log 2)^n} \simeq 3.25889 \frac{(n-1)!}{(2.07944)^n}. \quad (2)$$

*Proof of Theorem 1.* Take any Hamilton cycle  $C$ , of  $C_3(T_1, T_2, T_3)$ , and consider  $C$  restricted to  $T_1, T_2$ , and  $T_3$ . Since a Hamilton cycle meets every vertex in  $T_1, T_2$ , and  $T_3$  exactly once,  $C$  visits each subtournament the same number of times, say  $k$ . Hence for each  $T_i$ ,  $C$  will induce a collection of  $k$  disjoint paths that cover the vertices of  $T_i$ . We will refer to such a collection of  $k$  paths as a  $k$ -path cover. Similarly, given  $k$ -path covers for  $T_1, T_2, T_3$ , we can construct a Hamilton cycle by joining these  $k$ -path covers together. The number of ways of doing this is  $k!^3/k$ . Thus, if  $P(T_i, k)$  denotes the number of  $k$ -path covers of  $T_i$ , then the number of Hamilton cycles of  $C_3(T_1, T_2, T_3)$  which induce  $k$ -path covers in  $T_1, T_2$ , and  $T_3$  is

$$P(T_1, k)P(T_2, k)P(T_3, k) \frac{k!^3}{k}. \quad (3)$$

It follows that the number of Hamilton cycles in  $C_3(T_1, T_2, T_3)$  is

$$\sum_{k=1}^{\min\{m_1, m_2, m_3\}} P(T_1, k)P(T_2, k)P(T_3, k) \frac{k!^3}{k}.$$

For any set partition of the vertex set of  $T_i$  into  $k$  nonempty sets, each part will induce a subtournament of  $T_i$ . Rédei [5] showed that every tournament has a Hamilton path, thus each partition into  $k$  sets will induce at least one  $k$ -path cover of  $T_i$ . Therefore the number of Hamilton cycles in  $C_3(T_1, T_2, T_3)$  is at least

$$\sum_{k=1}^{\min\{m_1, m_2, m_3\}} S(m_1, k)S(m_2, k)S(m_3, k) \frac{k!^3}{k}$$

as claimed.

In the case that each  $T_i$  is transitive, each subtournament will have exactly one Hamilton path, hence we have equality in (1).  $\square$

*Proof of Corollary 2.* Suppose now that each  $T_i$  is a transitive tournament on  $m$  vertices, then the number of Hamilton cycles in  $C_3(T_1, T_2, T_3)$  is equal to

$$\sum_{k=1}^m S(m, k)^3 \frac{k!^3}{k}. \quad (4)$$

As with many combinatorial sums, the summands in (4) are approximated rather well by a normal distribution. Indeed, if we let

$$\mu = \frac{1}{2 \log 2} \quad \text{and} \quad \sigma = \frac{\sqrt{1 - \log 2}}{2 \log 2},$$

define  $f(m) = \sum_{k=1}^m S(m, k)k!$ , and write  $p(m, k) = \frac{S(m, k)k!}{f(m)}$ , then Bender [2] shows that  $p(m, k)$  is asymptotically normal with mean  $\mu m$  and variance  $\sigma^2 m$ . Hence,  $p(m, k)^3$  is also proportional to a normal distribution, at least in a range of  $k$  close to  $\mu m$ . This allows us to approximate the sum  $\sum_{k=1}^m S(m, k)^3 \frac{k!^3}{k}$  by an integral, showing that

$$\sum_{k=1}^m S(m, k)^3 \frac{k!^3}{k} \sim f(m)^3 \frac{3^{\frac{1}{2}} 2^{\frac{1}{2}} \pi^{\frac{1}{2}}}{3\mu\sigma^2 (2\pi)^{\frac{3}{2}} m^2}.$$

From Wilf [7, p. 176], we know that

$$f(m) \sim \frac{m!}{2(\log 2)^{m+1}},$$

Therefore with  $n = 3m$ , two applications of Stirling's approximation for  $n!$  yields

$$\begin{aligned} \sum_{k=1}^m S(m, k)^3 \frac{k!^3}{k} &\sim \left( \frac{m!}{2(\log 2)^{m+1}} \right)^3 \frac{3^{\frac{1}{2}} 2^{\frac{1}{2}} \pi^{\frac{1}{2}}}{3\mu\sigma^2 (2\pi)^{\frac{3}{2}} m^2} \\ &\sim \frac{\sqrt{2\pi n}}{n(1 - \log 2)} \left( \frac{n}{3e \log 2} \right)^n \\ &\sim \frac{1}{(1 - \log 2)} \frac{(n-1)!}{(3 \log 2)^n} \end{aligned}$$

□

**Acknowledgment.** The authors are very grateful to Rod Canfield for helpful advice regarding the asymptotics.

## References

- [1] N. Alon. The maximum number of Hamiltonian paths in tournaments *Combinatorica* (1990), 319-324.
- [2] E. A. Bender. Central and local limit theorems applied to asymptotic enumeration. *J. Combinatorial Theory Ser. A* 15 (1973), 91111.
- [3] B. Cuckler. Hamiltonian Cycles in Regular Tournaments *Combinatorics, Probability and Computing* (2007) 16, 239249.
- [4] J. W. Moon. *Topics on Tournaments* Holt, Reinhart and Winston (1968), New York.
- [5] L. Redei. Ein kombinatorischer Satz. *Acta Litt. Szeged*, 7:3943, 1934
- [6] T. Szele. Kombinatorikai vizsgálatok az irányított teljes graffal, *Kapcsolatban*, *Mt. Fiz. Lapok* 50 (1943), 223-256.

- [7] H. S. Wilf. Generatingfunctionology Academic Press, New York (1990).
- [8] N.C. Wormald,. Tournaments with many Hamilton cycles.