

10-1999

# Enhanced Biotransformation of Carbon Tetrachloride by *Acetobacterium woodii* upon Addition of Hydroxocobalamin and Fructose

Syed A. Hashsham  
*Michigan State University*

David L. Freedman  
*Clemson University*, [dfreedm@clemson.edu](mailto:dfreedm@clemson.edu)

Follow this and additional works at: [https://tigerprints.clemson.edu/envengineering\\_pubs](https://tigerprints.clemson.edu/envengineering_pubs)

 Part of the [Environmental Sciences Commons](#)

---

## Recommended Citation

Please use publisher's recommended citation.

This Article is brought to you for free and open access by the Environmental Engineering & Earth Sciences at TigerPrints. It has been accepted for inclusion in Publications by an authorized administrator of TigerPrints. For more information, please contact [kokeefe@clemson.edu](mailto:kokeefe@clemson.edu).

## Enhanced Biotransformation of Carbon Tetrachloride by *Acetobacterium woodii* upon Addition of Hydroxocobalamin and Fructose

SYED A. HASHSHAM<sup>1\*</sup> AND DAVID L. FREEDMAN<sup>2</sup>

Center for Microbial Ecology and Department of Civil and Environmental Engineering, Michigan State University, E. Lansing, Michigan 48824,<sup>1</sup> and Department of Environmental Engineering and Science, Clemson University, Clemson, South Carolina 29631<sup>2</sup>

Received 2 April 1999/Accepted 19 July 1999

The objective of this study was to evaluate the effect of hydroxocobalamin (OH-Cbl) on transformation of high concentrations of carbon tetrachloride (CT) by *Acetobacterium woodii* (ATCC 29683). Complete transformation of 470  $\mu\text{M}$  (72 mg/liter [aqueous]) CT was achieved by *A. woodii* within 2.5 days, when 10  $\mu\text{M}$  OH-Cbl was added along with 25.2 mM fructose. This was approximately 30 times faster than *A. woodii* cultures (live or autoclaved) and medium that did not receive OH-Cbl and 5 times faster than those controls that did receive OH-Cbl, but either live *A. woodii* or fructose was missing. CT transformation in treatments with only OH-Cbl was indicative of the important contribution of nonenzymatic reactions. Besides increasing the rate of CT transformation, addition of fructose and OH-Cbl to live cultures increased the percentage of [<sup>14</sup>C]CT transformed to <sup>14</sup>CO<sub>2</sub> (up to 31%) and <sup>14</sup>C-labeled soluble materials (principally L-lactate and acetate), while decreasing the percentage of CT reduced to chloroform and abiotically transformed to carbon disulfide. <sup>14</sup>CS<sub>2</sub> represented more than 35% of the [<sup>14</sup>C]CT in the presence of reduced medium and OH-Cbl. Conversion of CT to CO was a predominant pathway in formation of CO<sub>2</sub> in the presence of live cells and added fructose and OH-Cbl. These results indicate that the rate and distribution of products during cometabolic transformation of CT by *A. woodii* can be improved by the addition of fructose and OH-Cbl.

Metallocoenzymes like corrinoids, cytochrome P-450, and iron(II) porphyrins are known to play a major role in biotransformation and detoxification of carbon tetrachloride (CT) (22). The homoacetogen *Acetobacterium woodii* (1) is one of many organisms that transform CT at relatively high rates due in part to its use of the acetyl coenzyme A (CoA) pathway and correspondingly high levels of corrinoids (25). CT is transformed by fructose-grown *A. woodii* to mainly chloroform (CF), dichloromethane (DCM), and several soluble nonchlorinated products (9, 10, 24). Autoclaved *A. woodii* cultures also transform CT at an appreciable rate, perhaps indicating the heat stability of the enzymes involved. CT is also transformed by *Shewanella alga* BrY, an organism lacking the acetyl-CoA pathway (28). BrY reduces vitamin B<sub>12</sub>, which subsequently transforms CT primarily to carbon monoxide (CO).

These and other studies suggest that the role of microorganisms in dechlorinating CT may be limited to reduction of corrinoids or other cofactors, which can be accomplished equally well with chemical reducing agents (5, 17). However, previous results with a mixed culture grown on DCM and amended with cyanocobalamin indicates that an active microbial population may play a bigger role in CT transformation than simply providing reducing power (2, 14). Addition of cyanocobalamin significantly increases the rate of CT transformation and decreases CF accumulation, in comparison to autoclaved culture or sulfide-reduced basal medium supplemented with B<sub>12</sub>. Avoiding CF accumulation is a major concern with in situ bioremediation of CT.

The objective of this study was to determine the role of live

*A. woodii* cells in CT transformation when provided with fructose and hydroxocobalamin (OH-Cbl). Based on results from a methanogenic enrichment culture (14), we hypothesized that a pure culture of *A. woodii* supplemented with fructose and OH-Cbl would transform CT faster than autoclaved cells with the same amount of OH-Cbl and faster than OH-Cbl present in sulfide-reduced basal medium. In addition, we expected the presence of live cells supplemented with fructose and OH-Cbl to shift the product distribution from CT away from CF and DCM and towards CO and CO<sub>2</sub>. The results indicate that *A. woodii* recycles supplemental corrinoid to its reduced form, increases the rate of CT transformation to CO versus carbon disulfide (CS<sub>2</sub>), and drives metabolism of CO to CO<sub>2</sub> and fermentation products.

(Some preliminary results of this study were presented at the 96th Annual Meeting of the American Society for Microbiology, New Orleans, La., 19 to 23 May 1996.)

### MATERIALS AND METHODS

**Chemicals.** CT, CF, and DCM (all  $\geq 99.9\%$ ) and acetate, lactate, fumarate, and isobutyrate were obtained from Aldrich Chemical Co.; CS<sub>2</sub> was from EM Industries, Inc. OH-Cbl (98%) and cyanocobalamin (99%) were purchased from Sigma Chemical Co. [<sup>14</sup>C]CT was obtained from Dupont NEN Products and diluted to  $1.9 \times 10^7$  dpm/ml with distilled deionized water (0.54 mM CT). The radiochemical purity of this [<sup>14</sup>C]CT stock solution was evaluated to be  $99.6\% \pm 0.11\%$  ( $\pm 1$  standard deviation for duplicate bottles) by using the procedures described under "Analysis of <sup>14</sup>C products." EcoScint (Baker Diagnostics, Inc.) was used as liquid scintillation cocktail for all <sup>14</sup>C radioactivity measurements.

**Growth conditions.** *A. woodii* (ATCC 29683) was grown on fructose in a basal medium described by Freedman and Gossett (11), modified as follows: the Fe<sup>2+</sup> concentration was reduced to 1.2 mg/liter, and 10 ml of a vitamin solution per liter was added (27). Fructose was added as a sterile stock solution (2.8 or 25.2 mM).

*A. woodii* was grown in 160-ml serum bottles that were modified by connecting a 1-cm inside-diameter test tube at a right angle to the side of the bottles near the base, resulting in a final bottle volume of 173 ml. These modified serum bottles resemble culture flasks with a side arm (e.g., Bellco Biotechnology or Ace Glass), making it possible to monitor growth of *A. woodii* by optical density at 620 nm

\* Corresponding author. Present address: Department of Civil and Environmental Engineering, A126 Research Complex—Engineering, Michigan State University, East Lansing, MI 48824. Phone: (517) 355-8241. Fax: (517) 355-0250. E-mail: hashsham@pilot.msu.edu.

(Bausch and Lomb Spectronic 20 spectrophotometer). A correlation was developed between optical density and dry weight. Cells were harvested by centrifugation and resuspended in the basal medium. The suspension was filtered through a glass fiber filter (Whatman, 21 mm), dried at 105°C overnight, and weighed. An optical density of 0.1 using a 1-cm light path corresponded to 45 mg of cells per liter.

Basal medium was distributed to serum bottles in an anaerobic glove box (Coy Laboratory products, Inc.). The headspace of the bottles was then purged with an N<sub>2</sub>-CO<sub>2</sub> gas mixture (70%/30% [vol/vol]) to remove hydrogen and equilibrate the bicarbonate in the basal medium with CO<sub>2</sub> (resulting in a pH of 7.0). The serum bottles were sealed with slotted gray butyl rubber septa, covered with aluminum foil, and incubated at 35°C on a gyratory shaker table, with the liquid in contact with the septum.

**Volatile organic product analyses.** CT, CF, DCM, CH<sub>4</sub>, and CS<sub>2</sub> were measured by gas chromatographic analysis (Perkin-Elmer model 9000) of 0.5-ml samples taken from the 65-ml headspace of the serum bottles. A Carboxpak 1% SP-1000 column was used along with a flame ionization detector, as previously described (11). Although flame ionization is not the most suitable detector for measurement of CS<sub>2</sub>, a linear response was obtained for concentrations less than 50 µmol per bottle. Detection limits (nanomoles per bottle) were 60 for CH<sub>4</sub>, 20 for DCM, 100 for CS<sub>2</sub>, 20 for CF, and 4 for CT. CO was analyzed isothermally (90°C) on a gas chromatograph (GOW-MAC Instruments) equipped with a thermal conductivity detector (150°C), a 3.2-mm by 2.44-m stainless steel 80/100 mesh, Molecular Sieve 5A column (GOW-MAC Instruments), and helium as the carrier gas (30 ml/min). The detection limit for CO was approximately 1 µmol per bottle.

All of the CT transformation products reported in this study are in micromoles per bottle. Aqueous phase concentrations of the volatile compounds can be obtained by using their Henry's constant (mol·m<sup>-3</sup> gas concentration/mol·m<sup>-3</sup> aqueous concentration) at 35°C, as reported previously (14), and the liquid and headspace volumes in the serum bottles. The percentages of each compound found in the aqueous phase (108 ml) were 4.8% for CH<sub>4</sub>, 3.4% for CO, 93% for DCM, 56% for CS<sub>2</sub>, 88% for CF, and 47% for CT.

**CT transformation studies.** Experiments were initiated by adding 2 ml of a log-phase culture of *A. woodii* growing on fructose to 106 ml of basal medium in serum bottles. Initially 2.8 mM fructose was added, to grow the culture to a dry cell mass of 12.2 ± 0.44 mg/bottle. During this period, the bottles were not shaken. After growth was complete (i.e., no further increase in optical density) by day 3, 11 treatments were set up, using duplicate serum bottles for each treatment. CT (100 µmol per bottle), additional fructose (25.2 mM), and OH-Cbl (108 nmol per bottle) was then added to appropriate bottles, which were then monitored for optical density and CT transformation products. Treatments 1 to 7 consisted of sulfide-reduced basal medium (2.1 mM sulfide) with combinations of live or autoclaved *A. woodii*, OH-Cbl, and fructose. Water controls (treatment 8) consisted of 160-ml serum bottles with 100 ml of autoclaved deionized water and 100 µmol of CT, to evaluate losses through the septum and measure the radiochemical purity of the [<sup>14</sup>C]CT. Treatments 9 through 11 were included to check growth of *A. woodii* on fructose and/or OH-Cbl.

**Analysis of <sup>14</sup>C products.** [<sup>14</sup>C]CT was added (approximately 1.9 × 10<sup>6</sup> dpm) along with unlabeled CT to bottles representing treatments 1 to 8 (see Table 1). Treatment 1 (live *A. woodii* culture plus [<sup>14</sup>C]CT plus OH-Cbl plus fructose) included two identical sets; one analyzed for <sup>14</sup>C transformation products after 2.5 days and the other after 13 days (along with all of the other treatments) prior to analysis of <sup>14</sup>C products.

The amount of <sup>14</sup>C activity associated with volatile products was determined by gas chromatographic separation and combustion tube analysis of headspace samples (11). The efficiency of product recovery measured as trapped CO<sub>2</sub> after passing through the combustion tube was 95% ± 2.7% for CT, CF, and DCM and 98% ± 0.01% for methane. <sup>14</sup>CO<sub>2</sub> was determined by stripping acidified samples with N<sub>2</sub> and trapping in NaOH, as previously described (11).

Analysis of <sup>14</sup>C nonvolatile products was carried out by acidifying a 20-ml portion of the liquid sample with HCl and sparging with N<sub>2</sub> for 30 min. <sup>14</sup>C not purged at acid pH corresponded to nonstrippable residue (NSR). <sup>14</sup>C-labeled NSR retained by a 0.22-µm-pore-size filter (Whatman, Inc.) was presumed to be cell-associated. <sup>14</sup>C-labeled NSR filtrate (i.e., soluble NSR) was either concentrated 50- to 100-fold in a vacuum centrifuge before high-performance liquid chromatography (HPLC) analysis or injected directly onto the HPLC (10 to 250 µl). Separation was achieved on a 300-mm-diameter HPX-87H ion exclusion column (Bio-Rad Laboratories) connected to a UV-visible light absorbance detector (model 486; Millipore Corp.) set at 210 nm. Three different mobile phase conditions (0.013 N H<sub>2</sub>SO<sub>4</sub>, 0.7 ml/min, 30°C; 0.013 N H<sub>2</sub>SO<sub>4</sub> with 5% acetonitrile, 0.5 ml/min, 27°C; and 0.007 N H<sub>2</sub>SO<sub>4</sub> with 10.8% acetonitrile, 0.5 ml/min, 27°C) were used in order to confirm the identity of the compounds by shifts in retention times. The presence of acetate and butyrate was also confirmed by gas chromatographic analysis (Hewlett Packard 5890A series II) with a 4-mm inside diameter and 183-cm-long glass column (GP 10% SP-1200-1% H<sub>3</sub>PO<sub>4</sub> on 80/100 Chromosorb W AW; Supelco, Inc.).

To quantify which of the peaks from the HPX-87H column contained <sup>14</sup>C activity, fractions were collected in 0.5- to 1-min intervals, mixed with 15 ml of scintillation cocktail, and counted on a liquid scintillation counter. The identity of <sup>14</sup>C labeled L-lactate was also confirmed by its reactivity with lactate dehy-

drogenase. The average percent recovery during the HPLC analysis [ $\Sigma(^{14}\text{C}$  in all fractions)]/<sup>14</sup>C injected] was 89% ± 5.0%.

The recovery efficiency during the <sup>14</sup>C analyses was defined as the total disintegrations per minute recovered in all components (CT + CF + DCM + CS<sub>2</sub> + CO + CO<sub>2</sub> + pyruvate + L-lactate + acetate + isobutyrate + unidentified soluble NSR + cell associated fraction) divided by the total disintegrations per minute present at the time of analysis. The total disintegrations per minute present in a bottle at the time of <sup>14</sup>C analysis was the sum of 0.5-ml headspace sample disintegrations per minute and 100-µl aqueous phase sample disintegrations per minute measured by direct injection into scintillation cocktail. For bottles with low CS<sub>2</sub> levels (i.e., ≤22%), the recovery based on <sup>14</sup>C remaining was 96% ± 3.7%; for bottles with high CS<sub>2</sub> levels, the recovery was 80.8% ± 1.2%. All <sup>14</sup>C transformation products are expressed as percent of disintegrations per minute initially injected without correcting for losses through the septum during the incubation period. CT transformation rates are obtained by dividing the difference of initial and final aqueous CT concentrations by the number of days it took for the transformation, adjusted for losses in the water controls.

## RESULTS AND DISCUSSION

**Rate enhancement due to OH-Cbl addition.** The focus of this research was to determine if the rate of CT biotransformation by *A. woodii* can be enhanced by biochemical amendments, especially at high CT concentrations. OH-Cbl was selected for this purpose because cobalamins are one of the main coenzymes in the acetyl-CoA pathway implicated in the co-metabolic transformation of CT by *A. woodii* (10), and they are also known to transform CT in abiotic systems (18).

Results presented in Fig. 1a demonstrate that adding OH-Cbl and fructose to *A. woodii* caused a 30-fold increase in the rate of CT biotransformation with respect to controls that did not receive additional OH-Cbl (26 mg/liter/day for treatment 1 versus 0.83 mg/liter/day for treatment 5). When fructose or *A. woodii* cells were omitted or *A. woodii* cells were killed (treatments 2, 3, and 4), the addition of OH-Cbl still increased the rate fivefold, indicating the importance of abiotic sulfide-mediated CT transformations. The difference in rate between the treatment that contained *A. woodii* with fructose and OH-Cbl and all others that received only OH-Cbl indicates that *A. woodii* cells needed fructose to take advantage of the added OH-Cbl for CT transformation. The rate of CT transformation by *A. woodii* receiving only fructose was similar to previously reported rates (0.8 to 1.7 mg/liter/day) by Stromeyer et al. (24) and Egli et al. (10). Biomass concentrations estimated from protein and optical density data in all three studies were comparable (300 to 500 mg/liter). The amount of OH-Cbl (1 µmol/bottle) added in this study was only 1% of the initial CT, but it substantially increased the total amount of corrinoids present. Based on a corrinoid content of 0.52 nmol/mg of dry cell mass (7) and a cell mass of 40 mg/bottle, the corrinoid contributed by *A. woodii* cells was only 2% of the added OH-Cbl.

When OH-Cbl was replaced with an equimolar amount of cyanocobalamin in the live culture with fructose, there was only a marginal enhancement in the rate of CT transformation over live cells alone (data not shown), perhaps due to the toxicity of cyanide to *A. woodii* (20). Cyanocobalamin was successful in improving the rate of CT and CF transformation, as reported previously in a DCM-grown enrichment culture (2, 14) as well as in a mixed culture (15).

Since biomass concentration typically has a significant effect on the rate of CT transformation, growth was also monitored in all treatments involving *A. woodii*. The presence of CT inhibited growth of *A. woodii* (Fig. 1b). This was expected, since the aqueous CT concentration used was much higher than in most previous studies (72 versus <2 mg/liter). However, once CT was consumed and most of the accumulated CF was also transformed, (treatment 1; Fig. 1a and c), *A. woodii* grew to a cell density that was slightly higher than the maximum cell density for the controls with no CT. The absence of




FIG. 1. Transformation of CT (a), cell growth (b), and accumulation of chloroform (c), CO (d), and  $\text{CS}_2$  (e) by *A. woodii* and basal medium under various treatment conditions. 1, live *A. woodii* with CT, OH-Cbl, and fructose (●); 2, live *A. woodii* with CT and OH-Cbl but no fructose (◆); 3, autoclaved *A. woodii* with CT and OH-Cbl (■); 4, autoclaved medium with CT and OH-Cbl (▲); 5, live *A. woodii* with CT and fructose but no OH-Cbl (○); 6, autoclaved *A. woodii* with CT but no OH-Cbl (□); 7, autoclaved medium with CT but no OH-Cbl (△); 8, autoclaved water with CT but no OH-Cbl (×); 9, *A. woodii* seed control with fructose but no CT and no OH-Cbl (+); 10, *A. woodii* seed control with OH-Cbl and fructose but no CT (∇); and 11, *A. woodii* seed control with OH-Cbl but no fructose and no CT (▼). Vertical bars represent 1 standard deviation for duplicate bottles.

TABLE 1. Distribution of [<sup>14</sup>C]CT transformation products

Product	% of injected dpm by treatment <sup>a</sup>								
	1 <sup>b</sup>		2	3	4	5	6	7	8
	Live <i>A. woodii</i> + OH-Cbl (2.5 days)	Live <i>A. woodii</i> + OH-Cbl (13 days)	(live <i>A. woodii</i> + OH-Cbl, no fructose)	(autoclaved <i>A. woodii</i> + OH-Cbl)	(autoclaved medium + OH-Cbl)	(live <i>A. woodii</i> )	(autoclaved <i>A. woodii</i> )	(autoclaved medium)	(autoclaved water)
CT	0.62 ± 0.76	0	0	1.7 ± 0.41	0.52 ± 0.20	51 ± 0.29	48 ± 0.25	53 ± 0.28	65 ± 2.0
CF	3.4 ± 0.01	0	4.0 ± 0.29	4.1 ± 0.71	3.1 ± 0.21	4.1 ± 0.33	2.3 ± 0.04	0.88 ± 0.00	0
CS <sub>2</sub>	18 ± 1.3	13 ± 0.57	31 ± 1.8	30 ± 0.02	35 ± 0.12	1.9 ± 0.17	3.5 ± 0.15	2.1 ± 0.14	0
CO	25 ± 1.9	0.15 ± 0.06	8.6 ± 0.15	10 ± 0.05	6.5 ± 0.32	0.31 ± 0.21	0.77 ± 0.02	0	0
CO <sub>2</sub>	21 ± 0.44	31 ± 0.53	8.4 ± 0.38	5.0 ± 0.09	4.8 ± 0.64	2.4 ± 0.05	2.6 ± 0.23	1.6 ± 0.01	0.28 ± 0.06
Pyruvate	3.5 ± 0.09	0.98 ± 0.08	2.1 ± 0.19	1.5 ± 0.12	1.6 ± 0.14				
L-Lactate	2.5 ± 0.04	18 ± 0.31	1.8 ± 0.08	0	0				
Acetate	2.8 ± 0.13	6.5 ± 0.08	0.53 ± 0.08	1.9 ± 0.05	1.8 ± 0.11				
Isobutyrate	1.0 ± 0.03	3.6 ± 0.05	0.65 ± 0.05	0.93 ± 0.15	0.57 ± 0.05				
Other soluble NSR <sup>c</sup>	0.2 ± 0.04	1.3 ± 0.11	0.06 ± 0.02	0.31 ± 0.05	1.5 ± 0.07	2.1 ± 0.05	1.8 ± 0.15	1.1 ± 0.06	
Cell associated	4.3 ± 0.84	4.5 ± 0.28	2.2 ± 0.09	2.1 ± 0.04	1.4 ± 0.07	1.3 ± 0.08	0.95 ± 0.02	0.48 ± 0.05	

<sup>a</sup> Values represent the mean ± 1 standard deviation for duplicate bottles.

<sup>b</sup> Two sets in duplicate.

<sup>c</sup> For treatments 5 through 8, NSR was not analyzed for components. The soluble fraction of treatment 1 (live *A. woodii* plus OH-Cbl, 13 days) also included ethanol (2.3% ± 0.05%).

growth with fructose and OH-Cbl until nearly all of the CF was consumed (day 8, Fig. 1c) may be indicative of the activity of *A. woodii* towards toxicity reduction. There was no indication of OH-Cbl being used as a carbon source for growth in the presence or absence of fructose during the period of this study (Fig. 1b, treatments 9, 10, and 11).

#### Effect of OH-Cbl addition on volatile product distribution.

The distribution of CT transformation products is significantly influenced by experimental conditions, including the type of reducing environment, organisms present, and concentration of CT and coenzymes (5, 12, 16, 17, 26). The major intermediates observed during CT transformation are shown in Fig. 2. Egli et al. (10) presented a similar diagram based on studies with *A. woodii*, although it did not include CO or CS<sub>2</sub>. CO formation from CT is known to be catalyzed by corrinoids under reduced conditions. Further biotransformation of CO yields nonhazardous CO<sub>2</sub> and organic acids. CS<sub>2</sub> is produced from CT mainly under sulfide-mediated reducing conditions. Although it is a neurotoxin (13), its presence in drinking water is not currently regulated. CF is the most common undesirable transformation product of CT under various anaerobic conditions. DCM may also accumulate to some extent, via reduction of CF. Strategies that minimize accumulation of CF, DCM, and CS<sub>2</sub> are of interest for application of in situ bioremediation.

Addition of OH-Cbl promoted accumulation of CO (Fig. 1d, treatment 1). Without OH-Cbl, there was no detectable CO (below 1 μmol/bottle in treatments 5, 6, and 7). The increase in CO beyond day 8 corresponded to an increase in cells in bottles with *A. woodii*, OH-Cbl, and fructose (Fig. 1b), following depletion of the accumulated CF (Fig. 1c). Addition of OH-Cbl also promoted the accumulation of CS<sub>2</sub> (Fig. 1e). In the bottles with live culture plus OH-Cbl and fructose (treatment 1), the CS<sub>2</sub> peaked after 2.5 days and remained nearly constant thereafter, since all of the CT was consumed. CS<sub>2</sub> formation from CT has been observed in other microbial studies carried out in sulfide-reduced media (6, 14). However, neither Stromeyer et al. (24) nor Egli et al. (10) reported any CS<sub>2</sub> formation in their studies with *A. woodii* in a sulfide-reduced medium.

OH-Cbl addition also had a significant effect on accumulation of CF. In other studies with *A. woodii* (9, 10, 24) and

several anaerobic pure cultures (4, 19) without supplemental cobalamins, CF and DCM were among the major metabolites that accumulated during CT transformation. In this study also, accumulation of CF occurred in all of the treatments that received CT, but to a lower extent when OH-Cbl and fructose were also present with *A. woodii*.

The fastest rate of CT transformation and the highest level of CO accumulation occurred with the fructose and OH-Cbl amended live *A. woodii* (treatment 1), with a correspondingly lower level of CS<sub>2</sub> (compared to treatments 2, 3, and 4) and CF accumulation. Thus, active metabolism of an electron donor by *A. woodii* in the presence of supplemental OH-Cbl shifted the transformation of CT in favor of CO and away from CS<sub>2</sub> and CF. The combination of live cells, metabolism of the electron donor, and OH-Cbl was necessary to affect this shift. CT transformation still occurred at a high rate with sulfide-reduced OH-Cbl, but in the absence of fructose and active cells, CS<sub>2</sub> became the predominant product.

**Product distribution based on [<sup>14</sup>C]CT.** Use of [<sup>14</sup>C]CT made it possible to determine the distribution of soluble products and CO<sub>2</sub> originating from CT and to confirm the distribution of volatile products obtained by gas chromatographic analysis of headspace samples (Fig. 1). Table 1 shows the distribution of products from [<sup>14</sup>C]CT for each of the treatment conditions. The amount of [<sup>14</sup>C] lost during incubation was proportional to the total amount of CT remaining, as indicated by CT losses from the water controls (treatment 8). The maximum percentage of [<sup>14</sup>C]CO<sub>2</sub> we observed when fructose and OH-Cbl were added to live *A. woodii* cells (31%) was significantly higher than those reported in other pure culture studies, including *A. woodii* grown on fructose (9, 10, 24). Although oxidation of [<sup>14</sup>C]CO appears to be the principal mode of [<sup>14</sup>C]CO<sub>2</sub> formation in the presence of live cells, small amounts of [<sup>14</sup>C]CO<sub>2</sub> (<9%) did form with autoclaved cells and with basal medium alone (treatments 6 and 7); somewhat higher levels formed in the presence of supplemental OH-Cbl (treatments 3 and 4). How CT is converted to CO<sub>2</sub> in reduced medium without cells is not well established. One possibility is hydrolysis of CS<sub>2</sub> (17), which is favored at alkaline pH. With the OH-Cbl and fructose-supplemented active cultures, oxidation of CF may have contributed to CO<sub>2</sub> formation (2, 10). However, the extent of




FIG. 2. Pathways for biotransformation of CT by *A. woodii* in sulfide-reduced basal medium. Solid arrows represent major processes observed in this study; dashed arrows represent pathways reported elsewhere, but were minor in this study. Further reduction of DCM to chloromethane is possible, but occurs at very slow rates. Numbers in parentheses represent oxidation states.

CF accumulation was small relative to direct conversion of CT to CO. It should be noted that although CO was detected in treatment 1 by gas chromatographic analysis on day 13 (Fig. 1d), virtually none of it was  $^{14}\text{C}$  labeled at this point. The decrease in  $^{14}\text{CO}$  between days 2.5 and 13 in treatment 1 was accompanied by increases in  $^{14}\text{CO}_2$  and  $^{14}\text{C}$ -labeled L-lactate, acetate, ethanol, and isobutyrate. This is consistent with involvement of the acetyl-CoA pathway in CT transformation as proposed by Egli et al. (10); i.e., oxidation of CO to  $\text{CO}_2$  and formation of fermentation products. *A. woodii* typically produces 3 mol of acetate per mol of fructose under nonlimiting conditions, although formation of other compounds has been observed under stressed conditions (3). In this study, the stress was most likely due to CT and CF. Vitamin  $\text{B}_{12}$  and carbon monoxide dehydrogenase, two key enzymes in the acetyl-CoA pathway, are adversely affected by CT (8). This effect may have been partially mitigated by the addition of supplemental OH-Cbl.

The results of this study are in agreement with previous experiments that observed enhanced CT transformation by a methanogenic enrichment culture supplemented with cyanocobalamin (14). When provided with fructose and supplemental OH-Cbl, *A. woodii* plays a bigger role in CT transformation than simply supplying coenzymes that carry out nonenzymatic transformations. Metabolically active cells are needed to realize the highest transformation rates and most favorable product distribution. The mechanism by which this occurs needs further investigation. For example, it is not yet known if the added OH-Cbl acts extracellularly, (as shown with vitamin  $\text{B}_{12}$  and *Shewanella alga* BrY [28]), intracellularly, or by some combination of the two.

Supplemental addition of cobalamin along with an electron donor offers the prospect of accelerating the rate of in situ anaerobic transformation of CT in contaminated groundwater, as well as minimizing formation of hazardous metabolites like CF. The relatively high cost of cobalamins may be more than offset by a reduction in the time required for remediation, as well as minimization of hazardous products. One negative aspect of this approach is the formation of  $\text{CS}_2$  in the presence of sulfides, although it can be removed under aerobic conditions, if necessary (23). Adsorption of cobalamin to aquifer material has also been found to be low, with a retardation factor of 2.1

in sand (15). This is in the same range as for CT (21), suggesting that distribution of cobalamin along with an electron donor throughout a contaminated area is a feasible process.

#### ACKNOWLEDGMENTS

This research was supported in part by a grant from the Oak Ridge Institute for Science and Education, U.S. Department of Energy, and by the U.S. Army Construction Engineering Research Laboratory.

#### REFERENCES

- Balch, W. E., S. Schoberth, R. S. Tanner, and R. S. Wolfe. 1977. *Acetobacterium*, a new genus of hydrogen-oxidizing, carbon dioxide-reducing, anaerobic bacteria. *Int. J. Syst. Bacteriol.* **27**:355-361.
- Becker, J. G., and D. L. Freedman. 1994. Use of cyanocobalamin to enhance anaerobic biodegradation of chloroform. *Environ. Sci. Technol.* **28**:1942-1949.
- Buschhorn, H., P. Dürre, and G. Gottschalk. 1992. Purification and properties of the coenzyme A-linked acetaldehyde dehydrogenase of *A. woodii*. *Arch. Microbiol.* **158**:132-138.
- Castro, C. E., M. C. Helvenston, and N. O. Belser. 1994. Biodehalogenation, reductive dehalogenation by *Methanobacterium thermoautotrophicum*. Comparison with nickel(I)octaethylisobacteriochlorin anion. An F-430 model. *Environ. Toxicol. Chem.* **13**:429-433.
- Chiu, P. C., and M. Reinhard. 1995. Metallocoenzyme-mediated reductive transformation of carbon tetrachloride in titanium(III) citrate aqueous solution. *Environ. Sci. Technol.* **29**:595-603.
- Criddle, C. S., J. T. DeWitt, and P. L. McCarty. 1990. Reductive dehalogenation of carbon tetrachloride by *Escherichia coli* K-12. *Appl. Environ. Microbiol.* **56**:3247-3254.
- Dangel, W., H. Schulz, G. Diekert, H. König, and G. Fuchs. 1987. Occurrence of corrinoid-containing membrane proteins in anaerobic bacteria. *Arch. Microbiol.* **148**:52-56.
- Davidova, M. N., N. B. Tarasova, F. K. Mukhitova, and I. U. Karpilova. 1994. Carbon monoxide in metabolism of anaerobic bacteria. *Can. J. Microbiol.* **40**:417-425.
- Egli, C., S. Stromeyer, A. M. Cook, and T. Leisinger. 1990. Transformation of tetra- and trichloromethane to  $\text{CO}_2$  by anaerobic bacteria is a non-enzymic process. *FEMS Microbiol. Lett.* **68**:207-212.
- Egli, C., T. Tschan, R. Scholtz, A. M. Cook, and T. Leisinger. 1988. Transformation of tetrachloromethane to dichloromethane and carbon dioxide by *Acetobacterium woodii*. *Appl. Environ. Microbiol.* **54**:2819-2824.
- Freedman, D. L., and J. M. Gossett. 1991. Biodegradation of dichloromethane and its utilization as a growth substrate under methanogenic conditions. *Appl. Environ. Microbiol.* **57**:2847-2857.
- Gantzer, C. J., and L. P. Wackett. 1991. Reductive dechlorination catalyzed by bacterial transition-metal coenzymes. *Environ. Sci. Technol.* **25**:715-722.
- Graham, D. G., V. Amarnath, W. M. Valentine, S. J. Pyle, and D. C. Anthony. 1995. Pathogenic studies of hexane and carbon disulfide neurotoxicity. *Crit. Rev. Toxicol.* **25**:91-112.
- Hashsham, S., R. Scholze, and D. L. Freedman. 1995. Cobalamin-enhanced anaerobic biotransformation of carbon tetrachloride. *Environ. Sci. Technol.* **29**:2856-2863.
- Hashsham, S. A., and D. L. Freedman. 1997. Enhanced biotransformation of carbon tetrachloride by an anaerobic enrichment culture, p. 465-470. *In Proceedings of the Fourth International In Situ and On-Site Bioremediation Symposium*. New Orleans, La., 28 April to 1 May 1997. Battelle Press, Columbus, Ohio.
- Klecka, G. M., and S. J. Gonsior. 1984. Reductive dechlorination of chlorinated methanes and ethanes by reduced iron (II) porphyrins. *Chemosphere* **13**:391-402.
- Kriegman-King, M. R., and M. Reinhard. 1994. Transformation of carbon tetrachloride by pyrite in aqueous solution. *Environ. Sci. Technol.* **28**:692-700.
- Krone, U. E., R. K. Thauer, H. P. C. Hogenkamp, and K. Steinbach. 1991. Reductive formation of carbon monoxide from  $\text{CCl}_4$  and FREONs 11, 12, and 13 catalyzed by corrinoids. *Biochemistry* **30**:2713-2719.
- Picardal, F. W., R. G. Arnold, H. Couch, A. M. Little, and M. E. Smith. 1993. Involvement of cytochromes in the anaerobic biotransformation of tetrachloromethane by *Shewanella putrefaciens* 200. *Appl. Environ. Microbiol.* **59**:3763-3770.
- Ragsdale, S. W., L. G. Ljungdahl, and D. V. DerVartanian. 1983. Isolation of carbon monoxide dehydrogenase from *Acetobacterium woodii* and comparison of its properties with those of the *Clostridium thermoaceticum* enzyme. *J. Bacteriol.* **155**:1224-1237.
- Semprini, L., G. D. Hopkins, D. B. Janssen, M. Lang, P. V. Roberts, and P. L. McCarty. 1991. In-situ transformation of carbon tetrachloride under anoxic conditions. EPA/600/2-90/060. United States Environmental Protection Agency, Washington, D.C.
- Slater, T. F. 1984. Free-radical mechanism in tissue injury. *Biochem. J.* **222**:1-15.

23. **Smith, N. A., and D. P. Kelly.** 1988. Oxidation of carbon disulphide as the sole source of energy for the autotrophic growth of *Thiobacillus thioparus* strain TK-m. *J. Gen. Microbiol.* **134**:3041.
24. **Stromeyer, S. A., K. Stumpf, A. M. Cook, and T. Leisinger.** 1992. Anaerobic degradation of tetrachloromethane by *Acetobacterium woodii*: separation of dechlorinative activities in cell extracts and roles of vitamin B<sub>12</sub> and other factors. *Biodegradation* **3**:113–123.
25. **Stupperich, E., H. J. Eisinger, and B. Krautler.** 1988. Diversity of corrinoids in acetogenic bacteria: *p*-cresolylcobalamide from *Sporomusa ovata*, 5-methoxy-6-methylbezimidazolylcobamide from *Clostridium formicoaceticum* and vitamin B<sub>12</sub> from *Acetobacterium woodii*. *Eur. J. Biochem.* **172**:459–464.
26. **Ukrainczyk, L., M. Chibwe, T. J. Pinnavaia, and S. A. Boyd.** 1995. Reductive dechlorination of carbon tetrachloride in water catalyzed by mineral-supported biomimetic cobalt macrocycles. *Environ. Sci. Technol.* **29**:439–445.
27. **Wolin, E. A., M. J. Wolin, and R. S. Wolfe.** 1963. Formation of methane by bacterial extracts. *J. Biol. Chem.* **238**:2882–2886.
28. **Workman, D. J., S. L. Woods, Y. A. Gorbi, J. K. Fredrickson, and M. J. Truex.** 1997. Microbial reduction of vitamin B<sub>12</sub> by *Shewanella alga* strain BrY with subsequent transformation of carbon tetrachloride. *Environ. Sci. Technol.* **31**:2292–2297.