2001

Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina

William Conner

Clemson University, wconner@clemson.edu

Terrell Baker

New Mexico State University - Main Campus

B. Graeme Lockaby

Auburn University Main Campus

John A. Stanturf

USDA Forest Service

Marianne K. Burke

USDA Forest Service

Follow this and additional works at: https://tigerprints.clemson.edu/ag_pubs

Part of the [Forest Sciences Commons](#)

Recommended Citation

Please use publisher's recommended citation.

This Article is brought to you for free and open access by the Plant and Environmental Sciences at TigerPrints. It has been accepted for inclusion in Publications by an authorized administrator of TigerPrints. For more information, please contact kokeefe@clemson.edu.

SOUTH CAROLINA

Terrell T. Baker III *

College of Agriculture and Home Economics
New Mexico State University
Box 30003, MSC 3AE
Las Cruces, NM 88003-8003

William H. Conner
Baruch Forest Science Institute
P.O. Box 596
Georgetown, SC 29442

B. Graeme Lockaby
School of Forestry
Auburn University
108 M.W. Smith Hall
Auburn, AL 36849-5418

John A. Stanturf
Center for Bottomland Hardwoods Research
USDA Forest Service
P.O. Box 227
Stoneville, MS 38776

Marianne K. Burke
Center for Forested Wetlands Research
USDA Forest Service Southern Research Station
2730 Savannah Highway
Charleston, SC 29414

ACKNOWLEDGEMENTS

We would like to thank Westvaco Corporation for generously providing access to the study site under the sponsorship of the Coosawhatchie Bottomland Ecosystem Study. L. Wayne Inabinette provided invaluable field assistance throughout this study. We would also like to thank Marianne K. Burke for her support. Robin Clawson, Erik Schilling, Chuck Sykes, and Don Stoeckel provided field support. This study was funded by the US Forest Service Southern Research Station. We are appreciative of the suggestions provided by three anonymous reviewers.

Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina

ABSTRACT

The highly dynamic, fine-root component of forested wetland ecosystems has received inadequate attention in the literature. Characterizing fine root dynamics is a challenging endeavor in any system, but the difficulties are particularly evident in forested floodplains where frequent hydrologic fluctuations directly influence fine root dynamics. Fine root ($\leq 3\text{mm}$) biomass, production, and turnover were estimated for three soils exhibiting different drainage patterns within a mixed-oak community on the Coosawhatchie River floodplain, Jasper County, SC. Within a 45-cm deep vertical profile, 74% of total fine root biomass was restricted to the upper 15 cm of the soil surface. Fine root biomass decreased as the soil became less well-drained (e.g., fine root biomass in well-drained soil > intermediately drained soil > poorly drained soil). Fine root productivity was measured for one year using minirhizotrons and *in-situ* screens. Both methods suggested higher fine root production in better drained soils but showed frequent fluctuations in fine root growth and mortality, suggesting the need for frequent sampling at short intervals (e.g., monthly) to accurately assess fine root growth and turnover. Fine root production, estimated with *in-situ* screens, was 1.5, 1.8, and 0.9 Mg ha⁻¹ yr⁻¹ in the well-drained, intermediately drained, and poorly drained soils, respectively. Results from minirhizotrons indicated that fine roots in well-drained soils grew to greater depths while fine roots in poorly drained soils were restricted to surface soils. Minirhizotrons also revealed that the distribution of fine roots among morphological classes changed between well-drained and poorly drained soils.

INTRODUCTION

Productivity of forested wetland ecosystems has been the focus of numerous studies. Most commonly, productivity is estimated using aboveground parameters such as litterfall and stemwood production (Brinson et al., 1980; Conner and Day, 1992; Conner et al., 1993; Conner, 1994; Megonigal et al., 1997). Many investigators have acknowledged, however, that failure to include belowground data will seriously underestimate forest ecosystem productivity (Vogt et al., 1986b; Day and Megonigal, 1993). It has been suggested that fine root production accounts for up to 75 % of total net primary production (NPP) in some forests (Nadelhoffer and Raich, 1992). Similar to aboveground foliage, large amounts of fine roots die annually and can contribute a quantity of litter similar in magnitude to foliar litter (McClaugherty et al., 1984). Fine root dynamics, therefore, represent a significant source of energy and nutrient flow through forested systems, particularly for those systems that are subject to periodic disturbances that increase the frequency and extent of fine root turnover.

Forested wetlands are considered among the most dynamic of all forested ecosystems, and vegetation productivity within these systems has been addressed in many studies (Mitsch and Gosselink, 1993; Megonigal et al., 1997 among others). However, only a few investigations have characterized belowground productivity and the processes that contribute to fine root dynamics in forested wetlands (Powell and Day, 1991; Megonigal and Day, 1992; Day and Megonigal, 1993; Jones et al., 1996). Day and Megonigal (1993) suggested that omission of belowground data might cause previously accepted relationships between flooding and vegetation to be less accurate. Results from their study indicated that flooding reduced belowground allocation although aboveground production might remain similar across flooding regimes. Similarly, Brinson (1990) has summarized reports indicating that belowground

1 production may be much more sensitive to changes in soil oxidation - reduction potential than
2 aboveground production. The latter observation is in agreement with findings from upland
3 systems (Vogt et al., 1993) in relation to the highly responsive nature of fine roots to relatively
4 subtle changes in microenvironment. It is clear that to understand the critical productivity
5 function of forested wetlands, additional data on belowground production and the factors
6 controlling fine root dynamics are needed.

7 The lack of root data associated with studies of forested ecosystems is often noted (e.g.,
8 Vogt et al., 1986a; Megonigal et al., 1997; Lockaby and Walbridge, 1998). The aversion to
9 conducting root studies involves inherent difficulties associated with methodologies for studying
10 root systems. Most methods for estimating standing stocks of root biomass and/or production
11 involve three tasks: excavation, washing, and weighing (Caldwell and Virginia, 1989); the
12 former two are particularly labor intensive and time consuming. The common method for
13 estimation of root turnover requires sequential excavation to identify temporal fluctuations in
14 biomass that may be associated with production and mortality (Symbula and Day, 1988;
15 Caldwell and Virginia, 1989). A major challenge to this approach is identification of the
16 appropriate intervals at which to conduct sampling to accurately detect fluctuations in fine root
17 biomass (i.e., production and mortality). Vogt et al. (1986b) and Kurz and Kimmins (1987)
18 stressed that sampling be conducted at both peak and trough periods of fine root biomass to avoid
19 underestimation of production and mortality. Such timing is not as complicated for systems in
20 which fine root growth and mortality occur predictably. In the upland north-temperate hardwood
21 forest studied by Burke and Raynal (1994), for example, root growth was largely governed by
22 temperature. In southern floodplain forests, however, production and mortality are governed not
23 only by temperature but also by periodic flood events that occur at irregular and unpredictable

intervals. This prompts the need for more intensive sampling efforts at more frequent intervals – requiring considerably greater labor expenditure. The development of reliable sampling procedures that are less-labor intensive and time-consuming would be extremely helpful for characterizing belowground dynamics, particularly in southern forested floodplains.

While the processes controlling NPP in forested wetlands are complex, it is generally accepted that hydroperiod is the dominant controlling influence (Mitsch and Gosselink, 1993). There is disagreement, however, as to whether the flood events that are typical of forested wetlands represent a stress or a subsidy to vegetation in these systems (Conner and Day, 1976; Mitsch and Ewel, 1979; Megonigal et al., 1997). For example, Burke (*in press*) found continuously flooded stands were more productive than periodically flooded stands. In contrast, Megonigal et al.'s (1997) recent synthesis of studies characterizing productivity of numerous forested wetlands concluded that flooding tended to reduce aboveground NPP. Both suggested, however, that their conclusions considered only aboveground components, and that incorporation of belowground data would greatly improve our understanding of the productivity of entire wetland ecosystems.

It is important to consider not only the immediate effects of flooding on belowground productivity and turnover, but also the indirect effects resulting from many years of flood events, which shape floodplain landscapes and create a myriad of microsites within a single floodplain. It is common for an individual floodplain to exhibit a variety of soil microsites resulting from floodwater encroachment and recession (Jones et al., 1996). The vegetation mosaic created by the pattern of microsites within a single floodplain confounds the characterization of vegetation productivity, both above- and belowground, with each assemblage often exhibiting distinct production and allocation patterns. Often, such microsites differ in terms of soil chemistry, bulk

density, and, more dramatically, drainage characteristics. These characteristics in turn can play a significant role in fine root growth, production, and turnover.

The objectives of this study were to: 1) determine the vertical distribution of roots in three floodplain soils with different morphologies and drainage properties; 2) estimate and compare production of fine roots within each of these soils; and 3) examine the feasibility of two recent methods for estimating fine root production and phenology within a floodplain forest. Specifically, we hypothesized that: 1) most of the fine root biomass would be in the uppermost soil horizons; 2) fine root biomass would be lower in soils that were less well-drained; 3) fine roots would have a more shallow distribution in soils that were less well-drained; and 4) net fine root production would be reduced in soils that were less well-drained.

METHODS

STUDY SITE

This study was conducted in a bottomland oak community adjacent to the Coosawhatchie River, Jasper County, South Carolina on land owned by Westvaco Corporation (approximately 31° N, 81° W). Vegetation in the study area ranged from mixed oak at the higher portion of the study site to laurel oak-sweetgum-maple at the lower portion. The mixed oak stand had > 30% of the basal area in *Quercus phellos* L., *Q. nigra* L., and *Q. falcata* var. *pagodaefolia* Ell., with some *Pinus taeda* L. in the overstory. The laurel oak-sweetgum-maple stand had > 40% of the basal area in *Q. laurifolia* Michx., *Liquidambar styraciflua* L., and *Acer rubrum* L. These stand descriptions were based on a vegetation classification and ordination study and vegetation map by Burke and King (*in press*).

During the 1970's, USDA soil survey staff mapped the study area as a single unit; the Santee Association. A more intensive recent survey using 100 locations systematically located revealed and mapped nine distinct soil series throughout the floodplain (Murray et al. *in press*). The site at which our

study was located contained three soil series. According to the recent soil survey (Murray et al. *in press*), the highest part of the site was classified in the Coosaw series (thermic Arenic Hapludults) with silicious, sandy, and sandy loam surface layers exhibiting well-developed horizons and formed in older terrace sediments. The intermediate elevation at the site was in the Meggett series (fine, mixed, thermic Typic Albaqualfs). The lower, more poorly drained part of the site was classified in the Brookman series: fine, mixed, thermic Typic Umbraqualfs. These soils have thick, black loamy surface layers and dark gray clayey subsoils.

Preliminary observations of the study site revealed a tendency for floodwaters to remain above the soil surface for different lengths of time among the three soil series. The Coosaw series drained most rapidly followed by the Meggett series and the Brookman series, respectively. As some authors point out, surface flooding and hydroperiod represent only a fraction of actual hydrodynamics in floodplain ecosystems – the majority occurs below the soil surface and thus is not readily observable (Day et al., 1988; Day and Megonigal, 1993; Megonigal et al., 1997). Therefore, it was reasonable to assume that similar differences occurred belowground on this site and that these differences manifested themselves even in the absence of surface flooding. It was hypothesized that these differences were driven by differences in drainage conditions among the three soil series, and that this drainage gradient would be distinct even in the absence of flooding above the soil surface. The primary focus of this study was the difference in belowground production among the three soils resulting from this drainage differential. It should be recognized that the shift in vegetation, driven by the difference in drainage and water tolerance of the species present, as described above would have some effects on belowground processes such as fine root production, distribution, phenology, and nutrient dynamics. We make no attempt in this study to eliminate this source of variability.

SOIL CHARACTERISTICS

Five parallel transects (each 110 m long and 20 m apart) were installed and soil and fine root data were collected along each (Figure 1). The transects were installed across at least two of the soil series to test fine root response in relation to different soil drainage conditions. Soil temperature was monitored using six portable temperature recorders (Onset Computer Corporation, 1996), each of which was placed just below the soil surface, at 20-m intervals, along one of the transects. Steel welding rods were installed at the same locations to track monthly patterns of soil oxidation and Fe reduction (Bridgham et al., 1991). To complement welding rod measurements, soil coring was conducted to determine the depth to redoximorphic features (i.e., mottling, gleying). Bulk density measurements were taken in November 1995 using a 5-cm diameter bulk density probe to determine if the Coosaw, Meggett, and Brookman soil series differed in terms of bulk density. Exchangeable soil Ca, Mg, and K were determined on a Perkin-Elmer 373 Atomic Absorption Spectrophotometer following a double-acid extraction (Mehlich, 1953) and soil P was determined according to Watanabe and Olsen (1965). Soil pH was determined using a 1:1 soil:water ratio.

FINE ROOT DISTRIBUTION

The term "fine roots" is defined here as those roots having a diameter ≤ 3 mm. Although other studies have defined fine roots as being less than 2 mm, the 3-mm designation was chosen because a natural division seems to occur at approximately 3 mm as roots larger than this usually have secondary xylem thickening and tend to be perennial (McClagherty et al., 1982).

Fine root distribution was sampled along three of the transects, each containing 12 sample points, 10 m apart (Figure 1). At each sampling point, three soil cores, from each of three depths (0 - 15, 15 - 30, and 30 - 45 cm) were extracted using a 5-cm diameter bucket auger, for a total of 108 samples in March 1995. Sampling was confined to the top 45 cm of soil, as previous studies in similar systems have indicated that approximately 66% of fine roots may be restricted to that zone (Brown, 1990; Farrish,

1991). Samples were promptly placed in coolers, returned to the lab, and refrigerated at 4° C to maintain live roots until they could be analyzed (within 1 month). Soil cores were washed and sieved using a hydropneumatic root elutriator (Gillison's Variety Fabrication, Inc., Benzonia, MI). Root length was estimated using the line-intercept method (Newman, 1966) as described in Bohm (1979). After fine root length was determined, samples were oven-dried to a constant mass at 70° C, and dry mass was recorded for each depth.

FINE ROOT DYNAMICS AND PHENOLOGY

Two methods were employed to assess fine root phenology and growth - *in-situ* screens and minirhizotrons. While both methods have been used in upland systems (Fahey et al., 1989; Hendrick and Pregitzer, 1992), to our knowledge, their applicability in floodplain systems had not been determined.

In-Situ Screens

Melhuish and Lang (1968, 1971) have described a relationship between number of intersections that growing roots make with a plane of known area and estimated fine root length. In this study, six screens (*Phifer fiberglass 18/14 holes in²*, 7.6 cm x 15.2 cm), 1-m apart, were placed (using a sharpshooter or narrow, elongated spade) in the soil in April 1995 on a line perpendicular to the existing transect at each sampling point (Figure 1). Screens were oriented randomly with respect to aspect (i.e., N, S, E, or W) to prevent sampling bias in direction of root growth. Screens were inserted at 45° angles in the soil to correct for anisotropic root growth (Brown and Roussopoulos, 1974; Fahey and Hughes, 1994). One screen from each point along each transect was randomly selected for sampling during the first week of May, June, July, August, and November 1995 and April 1996. Screens were removed with a post hole digger and returned to the laboratory under refrigeration to be processed and analyzed as described above for soil core samples. Roots were separated from the screen and soil by hand as

mechanical techniques are not appropriate for this task. The following procedure was used to estimate fine root length and biomass production for each screen:

$$Eq. 1 \quad I_{cm2} = (I / SA)$$

$$Eq. 2 \quad L_{cm3} = 2 \times I_{cm2}$$

$$Eq. 3 \quad B = L_{cm3} \times b$$

$$Eq. 4 \quad G = B \times 107,700$$

where: I_{cm2} = # of root intersections cm^{-2} screen

I = # of intersections roots make with each screen

SA = surface area of a screen (cm^2)

L_{cm3} = root length cm^{-3} soil

2 = constant used to express length cm^{-2} of screen on volume basis (cm^{-3})*

B = biomass (g) of root cm^{-3} of soil*

b = biomass (g) of each cm of root*

G = root biomass (g) m^{-2} of soil surface to 10.77-cm depth

$107,700$ = expansion factor to achieve total biomass m^{-2} *

* see text for explanation

The number of intersections that fine roots made with each screen was counted (I). Based on the area sampled by each screen ($SA=116.13 \text{ cm}^2$), the number of intersections made with each cm^2 of the screen could be estimated (eq. 1). Melhuish and Lang (1968, 1971) demonstrated that random lines intersecting a cube, regardless of volume, would have a mean length of 0.6667 (the units corresponding to the volume of the cube under consideration). Multiplying this constant by the number of intersections that the lines make with one face of the cube (I_{cm2}), multiplying by six to consider all faces of the cube,

and dividing by 2 to account for each line intersecting the cube twice yields the equation $L_{cm3}=2 \times I_{cm2}$ (e.g., $L_{cm3} = \{6 \times 0.6667 \times I_{cm2}\}/2$). Therefore, Melhuish (1968) and Melhuish and Lang (1968, 1971) suggested that doubling the number of intersections that random lines make with one face of a cube (a plane) will accurately reflect the length of those lines within that cube. Their research using roots of cotton plants suggested that this approach would be useful for determining root length per unit volume of soil by simply examining one face (e.g., a plane or screen) of the cube. However, this approach assumes the following: 1) roots grow in a straight line for an infinitely short distance on either side of the plane; 2) roots are growing randomly in all angular directions; and 3) roots grow in all directions and at all places along the plane with equal probability (Melhuish and Lang, 1968).

In their studies, Melhuish (1968) and Melhuish and Lang (1968, 1971) discussed root length. While root length has been an important variable to consider, primarily in agronomic situations, root biomass is a preferred response variable for forested ecosystems as it enables interpretation of nutrient pools. Biomass per unit of volume of soil can be estimated by utilizing a simple length to biomass conversion. To accomplish this, a subsample of roots intersecting each screen at every sampling interval was taken. Roots were separated from the screens and length was estimated using the line-intercept method (Newman, 1966). Roots were then oven-dried to a constant mass and total mass was divided by total length for each sample to yield a constant for biomass per cm of root at each sampling period (b). Using the estimates of root length per unit volume of soil (L_{cm3}) derived from eq. 2, biomass per unit volume of soil was estimated using eq. 3. The expansion factor (107,700) expressed estimates on a square meter basis (eq. 4) (to an approximate depth of 10.7 cm) and was derived using the following: 100 cm x 100 cm x 10.77 cm (vertical depth of screens in soil).

Virtually no dead roots were observed with *in-situ* screen samples. It is likely that dead, and perhaps brittle, fine roots were lost during extraction of the screens. Mortality, therefore, was not directly measured but inferred from significant decreases in fine root biomass across sampling intervals.

1 Fine root N concentration was determined by thermal combustion using a Perkin-Elmer 2400
2 CHN-analyzer on subsamples taken from fine roots intersecting screens at each time period. Fine root P
3 concentration was determined colorimetrically using an ammonium vanadate solution (Jackson, 1958) on
4 a HCl extract following dry-ashing at 500° C for four hours. Fine root N and P contents were determined
5 by multiplying fine root N and P concentration by biomass as estimated with *in-situ* screens for each time
6 period.

7 8 Minirhizotrons

9 The minirhizotron technique was also used to monitor fine root dynamics on the study site. Six
10 clear acetate butyrate tubes (5-cm inside diameter, 5.7-cm outside diameter, 1.8-m long) were placed 20-
11 m apart on each of the two remaining transects, for a total of 12 tubes (Figure 1). Tubes were installed
12 during June 1995, when the water table was well below the soil surface. Installation of the tubes when
13 the soil was saturated may have resulted in inadequate seating and stability, thereby adversely
14 influencing root growth estimates. Tubes were inserted at 45° to limit the potential for roots to grow
15 along the soil/tube interface (Bragg et al., 1983). Aboveground portions of the tube were wrapped in
16 duct tape and capped to prevent entry of light and water. Care was taken to anchor tubes to ensure that
17 the fluctuating water table did not "push" tubes out of the soil. Standard metal conduit (5-cm diameter)
18 was hammered to a depth of approximately 1.5 m and anchored to each tube with a clamp and duct tape.

19 A fiberoptic periscope video camera (Bartz Technology, Inc. Santa Barbara, CA) was used to
20 monitor root growth once each month from August 1995 through July 1996. The camera was equipped
21 with a locking shaft to permit consistent, incremental lowering of the camera through tubes to a depth of
22 30 cm. Each time the camera was lowered, the number of root intersections with a predetermined grid
23 (the left and bottom sides of the monitor, in this case) was counted on the video monitor and recorded for
24 that depth. Roots were identified as live or dead based on their appearance and placed into one of three
25 morphological categories. Roots that appeared brown in color and exhibited characteristics of secondary

1 xylem thickening were classified as “Brown.” Roots that were translucent or white, and appeared
2 succulent were classified as “White.” Extremely small-diameter (< 0.5 mm) fine roots were classified as
3 “Hair-Like.” All roots classified as “Brown” or “White” were larger than 0.5 mm in diameter. Fine root
4 length was determined for the field counts using a variation of the line intercept method (Newman, 1966;
5 Bohm, 1979; Buckland et al. 1993). The Bartz camera was also equipped with an ultraviolet (UV) light
6 source that is designed to permit identification of live versus dead roots *in-situ*. According to Wang et al.
7 (1995), live roots will fluoresce when exposed to UV light. This feature is designed to eliminate
8 subjectivity in distinguishing live versus dead roots.

9 In addition to quantifying fine root length in the field using the video monitor, a video recording
10 was made one time each in August and September 1995 and every other month thereafter through July
11 1996. Recorded video images were then examined in the laboratory to test whether more intensive
12 analyses were feasible. Video images were digitized to computer using the Snappy Video Snapshot (Play
13 Incorporated, Rancho Cordova, CA, 1996), which connects to a parallel port on an IBM-compatible
14 personal computer. The Snappy Video Snapshot provided a low cost alternative (approximately \$100) to
15 expensive, hardware-intensive frame-grabbers. Each time the video camera was lowered, a new depth, or
16 field of view, was recorded on video tape. Each of these images was individually digitized as a frame on
17 which subsequent analyses could be conducted. On each frame, roots were classified as described above
18 for field counts. An image analysis software package (Optimas 6.0, Optimas Corporation, 1996) was
19 used to trace the perimeter of sufficiently large-diameter roots (“Brown” and “White”) and estimate their
20 diameter and planar surface area. Although automation is possible with this software, tracing was
21 conducted manually because the heterogeneity of the soil matrix in this system made automation
22 impractical. Due to their small diameter (e.g. < 0.5 mm), the perimeter of “Hair-Like” roots could not be
23 traced, rather a single line was drawn along the length of these structures. Calibrated according to the
24 area being analyzed (13.5 mm x 18 mm), Optimas returned the surface area of roots classified as either
25 “Brown” or “White” and the length of the “Hair-Like” roots. Roots classified as “Brown” or “White”

1 were treated as cylinders and the lengths of these structures were estimated using the diameter and
2 surface area estimates generated by Optimas ($\text{Length} = \text{Surface Area} / 2\pi r$). Tracing was conducted
3 manually because the heterogeneous soil background made automation difficult and subject to error.

4 5 STATISTICAL ANALYSES

6 Differences in fine root response variables (biomass, length, production) among soil series,
7 depth, and sample date were identified using the Student's t-test (PROC TTEST, SAS Institute, 1991).
8 Student's t-test was chosen rather than ANOVA because the study design was based on one experimental
9 unit for each soil series. Specifically, Student's t-tests were used within each soil series to compare
10 differences in fine root biomass and necromass among depths as well as differences within depths among
11 soil series. Also, production of fine root biomass was estimated using *in-situ* screens was compared
12 between soil series. Fine root length, as estimated using the minirhizotron method, was compared using
13 Student's t-test between soil series within each depth and fine root morphological category. Using the
14 same approach, fine root length among depths and morphological categories were also compared within
15 each soil series. Differences between means were considered statistically significant at $\alpha = 0.10$.
16 The less-conservative 90% level of significance was chosen due to the highly variable nature of fine root
17 data.

18 19 RESULTS

20 SOIL CHARACTERISTICS

21 Although bulk density and soil temperature data were collected during this study, no
22 differences in these variables were detected among drainage categories. Welding rod
23 measurements taken several times throughout the study indicated that the depth to reduced soil
24 conditions was lower in the Brookman series as compared to the Meggett series at every

sampling period (Table 1), and these differences were statistically significant in October 1995 and April 1996. Although welding rod data were not available for the Coosaw series, soil coring to determine the depth to mottling or gleying was conducted in May 1997 and confirmed that the depth to Fe reduction was greatest in this series. Each comparison between the Brookman series and the other two series was statistically significant, indicating less well-drained conditions in the Brookman series (Table 1). Results from both the welding rod measurements and the soil coring efforts confirmed that subsurface hydrology differed among the three soil series and revealed the hypothesized drainage differences which decreased in the order Coosaw, Meggett, and Brookman. Hereafter, these series will be discussed in terms of their drainage conditions: well-drained (WD), intermediately drained (ID), and poorly drained (PD), respectively. These terms are used as descriptors relative to each other and do not refer to any uniformly defined soil drainage categories or classifications.

Analysis of mineral elements in the three drainage categories indicated that the concentration of extractable P was greatest in the order: $WD > PD > ID$. Concentration of extractable K increased as soil drainage decreased such that $WD < ID < PD$. The patterns of extractable Ca and Mg were identical; concentration increased as successively less well-drained soils were encountered ($WD < ID < PD$, Table 1).

FINE ROOT DISTRIBUTION

In March 1995, the majority of fine roots (74 %) in this mixed-oak community were located in the upper 15 cm of soil as compared to 17 % and 9 % in the 15 - 30-cm and 30 - 45-cm depths, respectively (Figure 2). Fine root biomass tended to decrease with decreasing drainage

and depth (Figure 2). Because the majority of fine roots were located in the upper 15 cm of soil on the study site, subsequent efforts were directed toward fine roots in these surface soils.

FINE ROOT DYNAMICS AND PHENOLOGY

In-Situ Screens

Monthly estimates of fine root growth from *in-situ* screen samples are presented in Table 2. Statistical comparisons of fine root growth since installation of *in-situ* screens indicated few significant differences among the three drainage categories for each month. There was little difference between the WD and ID soils in terms of fine root production, and these differences were never statistically significant (Table 2). However, in April 1996, fine root production was significantly greater in the WD and ID soils as compared to production in the PD soil. Only in May 1995 did fine root production in the PD soil exhibit the greatest fine root production among the three drainage categories. In June 1995 fine root production in the PD soil was significantly greater than in the ID soil, but not in the WD soil.

Changes in nutrient content throughout the year are illustrated in Figure 3 and indicate the pool of each element contained in fine roots at each sampling interval. In terms of N content, the only significant differences among drainage categories were observed between the PD and ID soils. Only in June 1995 did the PD soil exhibit significantly higher fine root N content than the ID soil. The pattern observed was largely driven by the biomass of fine roots because few differences in N concentrations were observed among the drainage categories (data not shown, see Baker 1998). Exceptions to this occurred in June 1995 when fine root N concentration was significantly greater in the PD soil compared to the WD and ID soils and in November 1995 when fine root N concentration was significantly greater in the WD soil compared to the PD soil

(Baker 1998). Generally, fine root P content decreased such that $WD > ID > PD$, except during August 1995 (Figure 3). Similar to patterns observed for N content, differences in P content among drainage categories were driven largely by the biomass of roots sampled (Baker 1998). Ratios of N to P suggest subtle differences among the three drainage categories in terms of fine root litter quality but a consistent pattern across all sample dates was maintained such that ratios decreased in the order $PD > ID > WD$ (4.96, 3.54, and 2.60, respectively – data not shown).

Minirhizotrons

Field Counts

Field counts using the minirhizotron revealed a clear periodicity of fine root growth over the 12-month sampling period (Figure 4). Results are presented as root length density per unit of minirhizotron tube surface (mm cm^{-2}), similar to that reported by Day et al. (1996). However, our determination of root length differed in that we estimated root lengths based on the number of intersections with a grid on the screen in the field rather than through measurement of digitized images. Fine root length decreased from August 1995 to January 1996 with the exception of one short growth interval from October to November 1995. During the months of January 1996 through April 1996, fine root length fluctuated mildly but exhibited a brief increase from January to February 1996. As indicated on the May 1996 sample date, fine root growth accelerated during April 1996 and continued to increase in length until the final sample period in July 1996. With the exception of August and December 1995, the ID soil contained significantly higher root length density than the PD soil (Figure 4).

One of the advantages of using the minirhizotron method is the opportunity to examine root growth and morphological changes *in situ*. Relative changes in the proportion of each

morphological category can be monitored over time and with regard to particular environmental variables, which may influence the distribution of roots among morphological classes. For the purpose of illustration, Figure 5 compares the proportional distribution of fine roots encountered in the three morphological classes between the ID and PD soils. Within both drainage categories, the vast majority of fine root length was in the “Hair-Like” category. However, a greater proportion of fine roots in the PD soil fell into either the “White” or “Brown” classification.

Digitized Images

Similar to results from field counts, root length density across the entire depth sampled (0-30 cm) was statistically higher in the ID soil than in the PD soil for each sample period during which video images were recorded (data not shown). Preliminary analysis, as well as the results from the soil core samples collected in March 1995, suggested that frequency and distribution of fine roots began to decline beyond a certain depth. Therefore, the original 30-cm sampling depth was divided into two separate strata, 0 - 15 cm and 15 - 30 cm, for further analyses. Root length density was greater in the surface horizons than in the lower stratum for each drainage category, and these differences were statistically significant in all months except August 1995, for the PD soil (Table 3). Statistical analysis showed that, within the 0 - 15-cm depth, the ID soil contained greater root length density than the PD soil for every sample period and these differences were statistically significant except in May 1996 (Table 3). In the lower depth (15 - 30 cm), however, the PD soil contained higher root length density in August and September 1995. Apart from these two dates, the ID soil exhibited significantly higher root length density in the other months.

DISCUSSION

SOIL CHARACTERISTICS

Even though welding rod data were collected only intermittently for the ID and PD soils, and drainage data for the WD soil were based on soil cores, the hypothesized drainage pattern among the three soil series existed and was consistent such that $WD > ID > PD$, in terms of the depth to Fe reduction on the welding rods. It should be noted that measurements of the depth to Fe reduction on welding rods provide only an approximate estimate of water table fluctuations between sampling intervals and do not produce estimates as reliable as more intensive, well-monitoring data. Similarly, measurements based on soil morphological characteristics such as depth to mottling and gleying record longer-term hydrologic properties of soils.

All three soils exhibited low pH (Table 1). It is puzzling that soil-P concentration did not follow the gradient in drainage conditions (Table 1). Under successively waterlogged conditions, the concentration of P typically increases as Fe and Al complexes with P are reduced, thereby making the latter more available (Mitsch and Gosselink, 1993). Soil data in this study suggested the opposite trend, however, namely that P decreased as soil drainage decreased ($WD > PD > ID$, Table 1). It is probable that differences in the origin and genesis of each of the three soils, particularly the WD soil, accounted for the disparity in the observed pattern, but this was not tested.

FINE ROOT DISTRIBUTION

In a bottomland hardwood system in Louisiana, Farrish (1991) found that 64% of fine root biomass occurred in the top 20 cm of soil. Although Farrish (1991) sampled fine roots to a much greater depth, results from our study agree that the majority tend to be concentrated in the surface soils. Similarly, Symbula and Day (1988) and Powell and Day (1991) found greater fine

1 root biomass in surface soils than at lower depths in the Great Dismal Swamp. Fine roots are the
 2 structures primarily responsible for acquisition of water and nutrients (Marshall and Waring,
 3 1985; Farrish, 1991) and under conditions of comparatively low moisture, plants may allocate
 4 more resources to the construction of fine roots (Powell and Day, 1991; Marschner, 1995). The
 5 decrease in fine root biomass associated with poorer drainage observed in this study supports the
 6 hypothesis that fine root biomass increases along a gradient of decreasing soil moisture.
 7 However, it is unclear from our results whether the lower fine root biomass can be attributed to
 8 differences in vegetation across drainage categories, reduced growth in response to adequate
 9 moisture, hypoxia under poorly drained and thus poorly aerated conditions, or vegetation nutrient
 10 status.

12 FINE ROOT DYNAMICS AND PHENOLOGY

13 *In - Situ* Screens

14 Several points need to be made regarding the differences between the work described by
 15 Melhuish and Lang (1968, 1971) and its application here. Their earlier study was conducted on
 16 roots of cotton grown in a barrel of soil, and made the assumptions discussed above in the
 17 Methods section. The present study was conducted in a natural environment and may not adhere
 18 as stringently to the assumptions made by Melhuish and Lang (1968, 1971). However, it was
 19 assumed for this study that roots grew in all directions as well as angular directions. To capture
 20 growth in all directions, screens were oriented randomly throughout the study site. It was also
 21 assumed that roots grew at all places along the screen with equal probability, despite the fact that
 22 the majority of growth occurred in a fairly narrow band at the top of the screens. In two
 23 subsequent studies, Lang and Melhuish (1970) and Melhuish and Lang (1971) discussed the

1 implications for their technique in populations of roots that exhibit anisotropy and offered an
 2 anisotropy parameter to be used in calculating root length under those conditions. Because the
 3 degree to which roots were anisotropic could not be determined in the present study, this
 4 parameter was not used and our results are qualified by assuming that root growth is not only
 5 isotropic, but also fulfills the assumptions set forth in Melhuish and Lang's (1968, 1971) earlier
 6 studies.

7 Although several approaches for calculating production from changes in fine root
 8 biomass between sampling intervals have been discussed in the literature, only statistically
 9 significant increases in fine root standing stock between sample periods will be discussed in this
 10 study (Fairley and Alexander, 1985; Kurz and Kimmins, 1987; Symbula and Day, 1988;
 11 Publicover and Vogt, 1993). Several authors have contrasted this approach against simply
 12 summing all positive differences. For the purpose of comparison, we present estimates based on
 13 all positive increases between sampling intervals (Table 2).

14 Using only statistically significant increases in fine root biomass between sampling
 15 intervals, annual fine root NPP was 1,539, 1,810, and 937 kg ha⁻¹ yr⁻¹ to a depth of 10.77 cm for
 16 the WD, ID, and PD soils, respectively (Table 2). Inclusion of all increases, not just those that
 17 were statistically significant, would have resulted in considerably higher estimates of annual fine
 18 root NPP, and may have overestimated actual production. It is difficult to compare fine root
 19 production estimates reported in the literature. While most studies use similar diameter class
 20 designations, they vary in the use of a wide range of soil depths, time periods, and methods of
 21 calculating production. However, estimates from this study are within the ranges reported for
 22 other wetland systems. Jones et al. (1996) reported that fine root (≤ 5 mm) production to a depth
 23 of 20 cm ranged between 1030 and 6320 kg ha⁻¹ yr⁻¹ in a maple-gum community along a low-

order blackwater stream in Alabama using the sequential coring technique. Symbulla and Day (1988) reported higher fine root ($\leq 5\text{mm}$) production to a depth of 40 cm in a maple-gum community in the Great Dismal Swamp in Virginia; $5970\text{--}7830\text{ kg ha}^{-1}\text{ 11 mo}^{-1}$ and $6450\text{--}8860\text{ kg ha}^{-1}\text{ 11 mo}^{-1}$ using the implant bag and sequential coring techniques, respectively. Using the sequential coring technique, Powell and Day (1991) compared fine root ($\leq 5\text{mm}$) production to a depth of 40 cm between a mixed-hardwood community ($3540\text{--}9890\text{ kg ha}^{-1}\text{ yr}^{-1}$) and a maple-gum community ($590\text{--}910\text{ kg ha}^{-1}\text{ yr}^{-1}$).

Based on estimates derived from *in-situ* screens, fine roots experienced several pulses of growth and mortality throughout the year (Table 2). While it is possible that the decreases in fine root biomass observed in July 1995 and April 1996 could be the result of spatial variability in fine root biomass among sampling points, the fact that fine roots in all three drainage categories experienced mortality during the same months suggests that this was related to other edaphic factors. The decrease in biomass observed for the three drainage categories between June and July 1995 cannot be explained with the information presented here. However, these months are typically dry in this region and precipitation through July 1995 was 17.70 cm below normal based on historic data collected at the nearest National Oceanic and Atmospheric Administration (NOAA) station in Ridgeland, SC. Drought has been implicated in root mortality by earlier studies (Fogel, 1983). However, it has also been suggested that fine roots would respond to such conditions with increased growth (to a point) rather than the observed mortality (Keyes and Grier, 1981; Marschner, 1995).

In this floodplain community, it is unlikely that drought during the study was severe enough to significantly increase fine root mortality. Also, although contraction of heavy soils during extended dry spells has been shown to discourage fine root elongation (Marschner, 1995),

1 this does not explain the observed mortality as the *in-situ* screens were not installed deep enough
2 to contact soils with appreciable clay content (see soil descriptions in Methods). It has been
3 speculated that plants may respond to dry conditions by shifting root growth to greater depths
4 where water may be more abundant (Owensby et al., 1994). If this is true, the *in-situ* screens
5 may not receive new intersecting roots and may, in fact, lose fine roots as resources are allocated
6 to greater depths. It should be noted that virtually no dead fine roots were observed intersecting
7 *in-situ* screens in this study, perhaps due to loss during extraction or rapid decay. This method
8 did not, therefore, directly estimate fine root mortality – rather it was implied in the reduction of
9 quantities observed between sample periods.

10 Just as it is likely that fine root mortality during the dormant season accounted for some
11 of the decrease in observed fine root biomass in April 1996, it is also likely that poorly drained
12 conditions led to fine root mortality (Vogt et al., 1993; Marschner, 1995). Welding rod data for
13 April 1996 suggest that the water table was within 8 cm of the soil surface within the PD soil
14 during the preceding months (Table 1). Mortality under poor drainage is also supported by the
15 gradient of fine root biomass estimated with the *in-situ* screens: the WD soil maintained the
16 greatest biomass followed by the ID soil and finally the PD soil (Table 2). By far, the greatest
17 mortality was inferred in the PD soil.

18 Conversely, it also could be speculated that the relative position of the PD soil among the
19 drainage categories in terms of fine root growth in May and June 1995 may be the result of its
20 closer proximity to moisture during these dry months. Root growth that occurred in the PD soil
21 as a result of plants searching for moisture may have reversed the gradient of fine root growth
22 among the drainage categories from what was observed during wet months. In June 1996,
23 however, the WD soil exhibited the greatest fine root biomass and this phenomenon may be the

1 result of fine root growth exploiting a greater volume of soil for moisture uptake. Fine roots in
2 the ID and PD soils may not have responded similarly because conditions may not have been as
3 droughty in those soils.

4 These results indicate that fine roots in floodplain forests may experience several pulses
5 of production and mortality annually. This phenomenon suggests that studies of fine root
6 production and mortality must consider more intensive sampling intervals than would be
7 appropriate for other, less-dynamic systems. Approaches that measure fine root standing stocks
8 only twice each year may not reveal actual increases and decreases in fine root biomass and may,
9 therefore, seriously underestimate belowground production in floodplain systems (Vogt et al.,
10 1986b; Kurz and Kimmins, 1987). The maximum-minimum method for estimating fine root
11 production and mortality would not be appropriate in the mild climates of the southeastern
12 United States, particularly in floodplain forests where dynamic hydrologic processes contribute
13 substantially to the production and turnover of fine roots.

14 Although no significant patterns emerged with respect to fine root N contents determined
15 from samples intersecting *in-situ* screens, P content of fine roots appeared to be more sensitive to
16 soil drainage differences. Although P availability generally increases as soils become
17 progressively waterlogged and reduction of Fe and Al phosphates occurs (Mitsch and Gosselink,
18 1993), fine root P content in these soils do not appear to respond to this predicted P fertility
19 gradient. Results from this study suggested that P cycling through fine root turnover is greatest
20 along the drainage gradient in the order: WD > ID > PD.

21 Although blackwater rivers are usually associated with low primary productivity, net
22 productivity in this forested floodplain was among the highest reported for floodplain forests in
23 the South (Burke *in press*). This may be due, at least in part, to underlying marl deposited during

1 interglacial periods (Murray et al. *in press*) that contribute to the relatively high P and Ca
2 economy on the site. In a community that would normally be considered P-limited, the
3 Coosawhatchie site appears non-deficient in this element. It is not clear what effect, if any, this
4 may have on the patterns of fine root P concentration observed among the three drainage
5 categories under consideration.

6 Low ratios of N:P in fine roots that intersected *in-situ* screens suggested that this
7 floodplain was not P-limited. Generally N:P ratios > 15 suggest that the latter element is limiting
8 and microbial populations that utilize detritus will tend to immobilize P during decomposition
9 (Vogt et al., 1986a). On the nearby blackwater Ogeechee River in Georgia (approximately 100
10 km west), Lockaby et al. (1996) found that P was immobilized during decomposition of litterfall
11 exhibiting N:P ratios greater than 15. In the present study, N:P ratios of fine roots in all three
12 drainage categories remained well below this threshold value. It is interesting to note, however,
13 that fine root N:P ratios increased as drainage decreased.

14 15 Minirhizotrons

16 Ultraviolet illumination failed to allow us to distinguish between live and dead roots *in-*
17 *situ*. This problem has been identified in at least one other study (Wang et al., 1995). Despite
18 numerous field trials throughout the course of this investigation, ultraviolet light did not reliably
19 create fluorescence with roots that were known to be alive. Therefore, analyses that relied on
20 ultraviolet determination of live and dead roots were abandoned due to lack of confidence in the
21 procedure. Very few obviously dead roots were observed, and in most cases these were difficult
22 to distinguish from the soil matrix. An excellent review of the minirhizotron method,
23 particularly estimating root mortality, is presented in Hendrick and Pregitzer (1996).

Both the minirhizotron field counts and the digitizing procedures revealed seasonal fluctuations in root length density (Figure 4). Fine root growth and mortality were temporally similar to the patterns observed for roots sampled with *in-situ* screens (Table 2). Although water contamination and launch failures with temperature recorders precluded temperature estimates for some months, the trend illustrated in Figure 4 suggests that root elongation covaried with soil temperature.

Although minirhizotrons were installed only on the ID and PD soils, a pattern similar to that observed with *in-situ* screens was evident: the better-drained Meggett series maintained greater root length density than the poorly drained Brookman series. As several authors point out, it is not clear whether plants attempt to acquire resources (i.e. water and nutrients) by exploiting more thoroughly a given volume of soil or by exploring a greater volume of soil (Rogers et al., 1994; Day et al., 1996). This question is difficult to answer with the minirhizotron, as only small areas can be sampled. Results in this study suggested that fine roots tend to grow deeper during dry months, which supports the hypothesis that these roots explore a greater volume of soil in search of resources. Comparisons of the ID and PD soils suggested that roots in the ID soil exploited a greater soil volume than roots in the PD soil. It is unclear why the PD soil contained greater root length density of “White” and “Brown” roots than the ID soil. It would be expected that plants growing in the PD soil would be less likely to invest in more “permanent” structures given the tendency for these soils to become inundated.

Although similar patterns were obtained using both field counts and digitized images for gathering minirhizotron data, actual estimated root length densities differed between the two approaches. Despite good correlation (correlation coefficient, 0.78) between root length densities between field counts and digitized images, estimates from field counts were consistently higher

than results from analysis of digitized images. It is inevitable that the soil matrix contains inconsistencies and voids that present challenges to viewing roots at the tube/soil interface. During field sampling, the camera operator has the ability to use the focusing mechanism to improve the field of view. Once digitized, however, video images from minirhizotron samples are two-dimensional and the ability to improve the field of view is lost. This phenomenon may have contributed to the discrepancies between the two approaches and resulted in higher estimates for field counts. Other studies have reported good agreement between root lengths estimated from field counts and digitized images (e.g., $r^2=0.74$, Burch, 1995). That study was conducted in an upland system where less organic matter and lighter-colored, more homogenous soil would provide a better background against which roots could be observed during both procedures. This complication may have been exacerbated in this study by a frequently fluctuating water table that often obscured images and may have shifted soil materials around the minirhizotron tubes.

CONCLUSIONS

As we hypothesized, in a 45-cm soil profile within this floodplain oak community, most (74%) of the roots were restricted to the upper 15 cm of the soil. Our results also supported the hypothesis that fine root biomass would be lower in poorly drained soils as compared to more well-drained soils. Whereas well-drained soils contained higher fine root biomass in their surface depths as compared to poorly drained soils, the poorly drained soils contained a higher proportion of fine root biomass in their surface depths as compared to deeper strata. Fine root production was within the range reported in other bottomland hardwood studies. Although fine root NPP was greater in the well-drained and intermediately drained soils using only statistically

1 significant increases in biomass between sampling intervals, our results did not clearly support
2 the hypothesized decrease in fine root production with decreasing drainage. Although mortality
3 was not estimated directly, relative mortality inferred from *in-situ* screens and minirhizotrons
4 suggested that greater quantities of fine roots turn over annually in well-drained soils, despite the
5 fact that mortality appeared to be proportionally higher, and perhaps more frequent, in poorly
6 drained soils. In this landscape, fine roots in well-drained soils may contribute greater quantities
7 of higher-quality substrate to soil communities than poorly drained soils.

8 Both the minirhizotron and *in-situ* screen techniques revealed seasonal phenologies in
9 relation to soil temperature and, more significantly, soil drainage class. Both techniques appear
10 to be useful tools for monitoring fine root distribution and production and for estimating
11 mortality in frequently flooded, hydrologically dynamic floodplain ecosystems. Because these
12 methods are less time- and labor-intensive than traditional belowground sampling techniques,
13 they permit the more frequent sampling required in these systems. However, it should be
14 stressed that both techniques sample only small volumes of soil and are subject to the high spatial
15 and temporal variability inherent in fine root measurement. Therefore, increasing the number of
16 samples taken or points measured should be considered during their use. The application of
17 these techniques may be most useful for making comparisons among treatments as there is, as
18 yet, no reliable standard with which to compare actual production and mortality estimates
19 (Hendricks et al., 1993).

LITERATURE CITED

- 1
2 Baker, T. T. 1998. Fine Root Dynamics on a Forested Floodplain and Litter Decomposition in
3 Four Forested Floodplain Communities in the Southern United States. Ph.D.
4 Dissertation. Auburn University. Auburn, Alabama. 118 pp.
5
- 6 Bohm, W. 1979. Methods of Studying Root Systems. Springer-Verlag, New York. 188 p.
7
- 8 Bragg, P.L., G. Govi, and R.Q. Cannel. 1983. A comparison of methods, including angled and
9 vertical minirhizotrons, for studying root growth and distribution in spring oat crop.
10 Plant Soil 73: 435-440.
11
- 12 Bridgham, S. D., S. P. Faulkner, and C. J. Richardson. 1991. Steel rod oxidation as a hydrologic
13 indicator in wetland soils. Soil Sci. Soc. Am. J. 55: 856-862.
14
- 15 Brinson, M. 1990. Riverine forests. Pages 87-134 In: A. Lugo, M. Brinson, and S. Brown (Eds.).
16 Forested Wetlands. Ecosystems of the World 15. Elsevier Scientific Publishing,
17 Amsterdam.
18
- 19 Brinson, M. M., H. D. Bradshaw, R. N. Holmes, and J. B. Elkins, Jr. 1980. Litterfall, stemflow,
20 and throughfall nutrient fluxes in an alluvial swamp forest. Ecology 61: 827-835.
21
- 22 Brown, J. K. and P. S. Roussopoulos. 1974. Eliminating bias in the planar intersect method for
23 estimating volumes of small fuels. For. Sci. 20: 350-356.

- 1 Brown, S. 1990. Structure and dynamics of basin forested wetlands in North America. Pages
2 171-196 In: A. Lugo, M. Brinson, and S. Brown (Eds.). *Forested Wetlands. Ecosystems*
3 *of the World* 15. Elsevier Scientific Publishing, Amsterdam.
- 4
- 5 Buckland, S. T., C. D. Campbell, L. A. Mackie-Dawson, G. W. Horgan, and E. I. Duff. 1993. A
6 method for counting roots observed in minirhizotrons and their theoretical conversion to
7 root length density. *Plant Soil* 153: 1-9.
- 8
- 9 Burch, W.H. 1995. Rooting habits of loblolly pine, hardwoods, and herbaceous plants during
10 forest regeneration: a minirhizotron study. MS Thesis. Auburn University. Auburn, AL.
11 76 pp.
- 12
- 13 Burke, M.K. and D.J. Raynal. 1994. Fine root growth phenology, production, and turnover in a
14 northern hardwood forest ecosystem. *Plant Soil* 162: 135-146.
- 15
- 16 Burke, M.K., Liechty, H.O., and M.H. Eisenbies. 2000. Aboveground and belowground net
17 primary production. Pages 39-41 In: M.K. Burke and M.H. Eisenbies (Eds.). *The*
18 *Coosawhatchie Bottomland Ecosystem Study: a report on the development of a reference*
19 *wetland*. USDA Forest Service Southern Research Station, Asheville, North Carolina,
20 Gen. Tech. Rep. SRS-38.
- 21
- 22 Burke, M.K. King, S.L., Eisenbies, M.H. and D. Gartner. 2000. Vegetation and Soils. Pages
23 23-28 In: M.K. Burke and M.H. Eisenbies (Eds.). *The Coosawhatchie Bottomland*

1 Ecosystem Study: a report on the development of a reference wetland. USDA Forest
2 Service Southern Research Station, Asheville, North Carolina, Gen. Tech. Rep. SRS-38.

3
4 Caldwell, M.M. and R.A. Virginia. 1989. Root systems. Pages 367-398 In: R.W. Pearcy, J.R.
5 Ehleringer, H.A. Mooney, and P.W. Rundel (Eds.). Plant Physiological Ecology,
6 Chapman and Hall, London.

7
8 Conner, W. H. 1994. Effect of forest management practices on southern forested wetland
9 productivity. Wetlands 14: 27-40.

10
11 Conner, W. H. and J. W. Day, Jr. 1976. Productivity and composition of a baldcypress-water
12 tupelo site and bottomland hardwood site in a Louisiana swamp. Am. J. Bot. 63: 1354-
13 1364.

14
15 Conner, W. H. and J. W. Day, Jr. 1992. Water level variability and litterfall productivity of
16 forested freshwater wetlands in Louisiana. Am. Midl. Nat. 128: 237-245.

17
18 Conner, W. H., J. W. Day, Jr. and W. R. Slater. 1993. Bottomland hardwood productivity: case
19 study in a rapidly subsiding Louisiana, USA, watershed. Wetlands Ecology and
20 Management 2: 189-197.

21

Day, F. P. and J. P. Megonigal. 1993. The relationship between variable hydroperiod, production allocation, and belowground organic matter turnover in forested wetlands. *Wetlands* 13(2): 115-121.

Day, F. P., S. K. West, and E. G. Tupacz. 1988. The influence of groundwater dynamics in a periodically flooded ecosystem, the Great Dismal Swamp. *Wetlands* 8: 1-13.

Day, F. P., E. P. Weber, C. R. Hinkle, and B. G. Drake. 1996. Effects of elevated atmospheric CO₂ on fine root length and distribution in an oak-palmetto scrub ecosystem. *Global Change Biology* 2: 143-148.

Fahey, T. J. and J. W. Hughes. 1994. Fine root dynamics in a northern hardwood forest ecosystem, Hubbard Brook Experimental Forest, NH. *J. Ecol.* 82: 533-548.

Fahey, T.J., X. Zhihua, and J.W. Hughes. 1989. Fine root growth and turnover at Hubbard Brook Experimental Forest, NH. *Bull. Ecol. Soc. Am.* 70: 109-115.

Fairley, R. I. and I. J. Alexander. 1985. Methods of calculating fine root production in forests. Pages 37-42 In: A. H. Fitter, D. Atkinson, D. J. Read, and M. B. Usher, (Eds.). *Ecological Interactions in Soil*. Blackwell Scientific Publications, Boston, MA.

Farrish, K. W. 1991. Spatial and temporal fine-root distribution in three Louisiana forest soils. *Soil Sci. Soc. Am J.* 55: 1752-1757.

1 Fogel, R. 1983. Root turnover and productivity of coniferous forests. *Plant Soil* 71: 75-85.

2
3 Hendrick, R. L. and K. S. Pregitzer. 1996. Applications of minirhizotrons to understanding root
4 function in forests and other natural ecosystems. *Plant Soil* 185: 293-304.

5
6 Hendrick, R. L. and K. S. Pregitzer. 1992. The demography of fine roots in a northern
7 hardwood forest. *Ecology* 73(3): 1094-1104.

8
9 Hendricks, J. J., K. J. Nadelhoffer, and J. D. Aber. 1993. Assessing the role of fine roots in
10 carbon and nutrient cycling. *Tree* 8(5): 174-178.

11
12 Jackson, M. L. 1958. *Soil Chemical Analysis*. Prentice-Hall. Englewood Cliffs, New Jersey.
13 498 p.

14
15 Jones, R. H., B. G. Lockaby, and G. L. Somers. 1996. Effects of microtopography and
16 disturbance on fine-root dynamics in wetland forests of low-order stream floodplains.
17 *Am. Midl. Nat.* 136: 57-71.

18
19 Keyes, M. R. and C. C. Grier. 1981. Above- and below-ground net production in 40-yr-old
20 Douglas-fir stands on low and high productivity sites. *Can. J. For. Res.* 11: 599-605.

1 Kurz, W. A. and J. P. Kimmins. 1987. Analysis of some sources of error in methods used to
2 determine fine root production in forest ecosystems: a simulation approach. *Can. J. For.*
3 *Res.* 17: 909-912.

4
5 Lang, A. R. G. and F. M. Melhuish. 1970. Lengths and diameters of plant roots in non-random
6 populations by analysis of plane surfaces. *Biometrics* 26(3): 421-431.

7
8 Lockaby, B. G. and M. R. Walbridge. 1998. Biogeochemistry. Pages 149-172 In: M. G.
9 Messina and W. H. Conner (Eds.). *Southern Forested Wetlands: Ecology and*
10 *Management*. Lewis Publishers, Boca Raton, Florida.

11
12 Lockaby, B. G., A. L. Murphy, and G. L. Somers. 1996. Hydroperiod influences on nutrient
13 dynamics in decomposing litter of a floodplain forest. *Soil Sci. Soc. Am. J.* 60: 1267-
14 1272.

15
16 Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. Second Edition. Academic Press Inc.,
17 San Diego, California. 889 pp.

18
19 Marshall, J. D. and R. H. Waring. 1985. Predicting fine root production and turnover by
20 monitoring root starch and soil temperature. *Can. J. For. Res.* 15: 791-800.

21
22 McClaugherty, C.A., J.D. Aber, and J.M. Melillo. 1982. The role of fine roots in the organic
23 matter and nitrogen budgets of two forested ecosystems. *Ecology* 63(5): 1481-1490.

1 McClaugherty, C.A., J.D. Aber, and J.M. Melillo. 1984. Decomposition dynamics of fine roots
2 in forested ecosystems. *Oikos* 42: 378-386.

4 Megonigal, J. P. and F. P. Day. 1992. Effects of flooding on root and shoot production of bald
5 cypress in large experimental enclosures. *Ecology* 73(4): 1182-1193.

7 Megonigal, J. P., W. H. Conner, S. Kroeger, and R. R. Sharitz. 1997. Aboveground production
8 in southeastern floodplain forests: a test of the subsidy-stress hypothesis. *Ecology* 78(2):
9 370-384.

11 Mehlich, A. 1953. Determination of calcium, magnesium, potassium, sodium, and ammonium
12 by North Carolina soil testing laboratories. North Carolina State University Mimeograph,
13 Raleigh, North Carolina.

15 Melhuish, F. M. 1968. A precise technique for measurement of roots and root distribution in
16 soils. *Annals of Botany* 32: 15-22.

18 Melhuish, F. M. and A. R. G. Lang. 1968. Quantitative studies of roots in soil. 1. Lengths and
19 diameters of cotton roots in a clay-loam soil by analysis of surface-ground blocks of
20 resin-impregnated soil. *Soil Sci.* 106(1): 16-22.

22 Melhuish, F. M. and A. R. G. Lang. 1971. Quantitative studies of roots in soil. 2. Analysis of
23 non-random populations. *Soil Sci.* 112(3): 161-166.

1 Mitsch, W.J. and K.C. Ewel. 1979. Comparative biomass and growth of cypress in Florida
2 wetlands. *Am. Midl. Nat.* 101: 417-426.

3
4 Mitsch, W.J. and J.G. Gosselink. 1993. *Wetlands*, 2nd edition. Van Nostrand Reinhold, New
5 York, New York.

6
7 Murray, L.A., Eppinette, B., and J.H. Thorpe. 2000. Geomorphology and soil survey. Pages 7-9
8 In: M. K. Burke and M.H. Eisenbies (Eds.), *The Coosawhatchie Bottomland Ecosystem*
9 *Study: a report on the development of a reference wetland*. USDA Forest Service
10 Southern Research Station, Asheville, North Carolina, Gen. Tech. Rep. SRS-38.

11
12 Nadelhoffer, K.J. and J.W. Raich. 1992. Fine root production estimates and belowground
13 carbon allocation in forest ecosystems. *Ecology* 73(4): 1139-1147.

14
15 Newman, E. I. 1966. A method of measuring the total length of root in a sample. *J. Appl. Ecol.*
16 3: 139-145.

17
18 Owensby, C. E., L. M. Auen, and P. I. Coyn. 1994. Biomass production in a nitrogen-fertilized,
19 tallgrass prairie ecosystem exposed to ambient and elevated levels of CO₂. *Plant Soil*
20 165: 105-113.

21
22 Powell, S. W. and F. P. Day. 1991. Root production in four communities in the Great Dismal
23 Swamp. *Am. J. Bot.* 78: 288--297.

- Publicover, D. A. and K. A. Vogt. 1993. A comparison of methods for estimating forest fine root production with respect to sources of error. *Can. J. For. Res.* 23: 1179-1186.
- Rogers, H. H., Runion, G.B. and S. V. Krupa. 1994. Plant responses to atmospheric CO₂ enrichment of cotton: vertical and lateral root distribution patterns. *Plant Soil* 165: 33-44.
- SAS Institute. 1991. *SAS User's Guide: Statistics*. SAS Institute, Inc., Cary, North Carolina.
- Symbula, M. and F.P. Day. 1988. Evaluation of two methods for estimating belowground production in a freshwater swamp forest. *Am. Midl. Nat.* 120(2): 405-415.
- Vogt, K.A., C.C. Grier, and D.J. Vogt. 1986a. Production, turnover, and nutrient dynamics of above- and belowground detritus of world forests. *Adv. Ecol. Res.* 15: 303-377.
- Vogt, K.A., C. C. Grier, S. T. Gower, D. C. Sprugel. 1986b. Overestimation of net root production: a real or imaginary problem? *Ecology* 67: 577-579.
- Vogt, K.A., D.A. Publicover, J. Bloomfield, J.M. Perez, D.J. Vogt, and W.L. Silver. 1993. Belowground responses as indicators of environmental change. *Environ. Exp. Bot.* 33(1): 189-205.

1 Wang, Z., W.H. Burch, P. Mou, R.H. Jones, and R.J. Mitchell. 1995. Accuracy of visible and
2 ultraviolet light for estimating live root proportions with minirhizotrons. *Ecology* 76(7):
3 2330-2334.

4

5 Watanabe, FS and SR Olsen. 1965. Test of an ascorbic acid method for determining phosphorus
6 in water and NaHCO₃ extracts from soil. *Soil Sci. Soc. Amer. Proc.* 29: 677-678.

7

Table 1 Descriptive characteristics of a well-drained (WD), an intermediately drained (ID), and a poorly drained (PD) soil within a mixed-oak community on the Coosawhatchie River floodplain, Jasper County, SC. Standard errors of the means are in parentheses.

Depth to Reduction [†] (cm)			
Date	WD Coosaw Series	ID Meggett Series	PD Brookman Series
Aug '95	56 (7) a	64 (9) a [‡]	48 (5) a
Oct '95		19 (7) a	8 (1) b
Nov '95		37 (6) a	17 (3) a
Apr '96		24 (6) a	8 (1) b
Jun '96		55 (9) a	22 (5) a
Depth to Mottling/Gleying (cm)			
Date	WD Coosaw Series	ID Meggett Series	PD Brookman Series
May '97	56 (7) a	21 (5) a	3 (3) b
Chemistry (mg kg ⁻¹) [§]			
Parameter	WD Coosaw Series (0 - 18 cm)	ID Meggett Series (0 - 15 cm)	PD Brookman Series (0 - 18 cm)
pH	4.5	4.4	4.7
P [¶]	30.06	8.18	10.10
K	31.20	58.50	89.70
Mg	19.20	43.20	189.60
Ca	54.00	86.00	960.00

[†] Measured with steel welding rods.

[‡] Means with same lowercase letter in a row are not statistically different (alpha = 0.10).

[§] Murray et al. (*in press*).

[¶] This study.

Table 2 Monthly and annual fine root production estimates (kg ha^{-1}) for a well-drained (WD), an intermediately drained (ID), and a poorly drained (PD) soil, to a depth of 11 cm, as measured with *in-situ* screens for a mixed-oak community on the Coosawhatchie River floodplain, Jasper County, SC. Fine root growth was assumed to be zero at the time *in-situ* screens were installed.

	WD	Monthly NPP [†]	ID	Monthly NPP	PD	Monthly NPP
May '95	84.10 a [§]	84.10 [‡] **	91.83 a	91.83 **	115.76 b	115.76 **
June '95	941.18 ab	857.08 **	808.92 a	717.09 **	937.12 b	821.36 **
July '95	604.56 a	-336.62	603.99 a	-204.93	533.43 a	-403.69
Aug '95	1202.54 a	597.98 **	1604.99 a	1001.00**	700.11 a	166.68
Nov '95	1736.24 a	533.7	1608.99 a	4.00	1510.16 a	810.05
April '96	1624.45 b	-111.79	1529.73 b	-79.26	793.46 a	-716.70
Fine Root NPP (yr^{-1}) [¶]		2072.86		1813.92		1913.85
Fine Root NPP (yr^{-1}) [#]		1539.16		1809.92		937.12

[†] NPP represents net primary production.

[‡] An ** indicates statistically significant positive net increases in fine roots over previous month.

[§] Lowercase letters compare means between drainage categories for each month.

[¶] Calculation of annual fine root NPP by summing increases in fine roots over the previous month.

[#] Calculation of annual fine root NPP using only statistically significant increases in fine roots over previous month (Fairley and Alexander 1985, Kurz and Kimmins 1987, Publicover and Vogt 1993).

Table 3 Statistical comparison of root length density (mm cm^{-2}) between an intermediately drained (ID) and a poorly drained (PD) soil within two depth strata using digitized images from minirhizotron sampling, Coosawhatchie River floodplain, Jasper County, SC. Standard errors of the means are in parentheses.

Date	0 - 15 cm			15 - 30 cm		
	ID	PD	P > F	ID	PD	P > F
Aug '95	5.63 (0.89)	3.11 [†] (0.37)	0.001	1.76 (0.33)	1.92 [†] (0.36)	0.001
Sept '95	4.92 (0.93)	2.37 (0.31)	0.001	1.20 (0.32)	1.35 (0.27)	0.001
Nov '95	5.47 (0.96)	3.19 (0.51)	0.001	1.09 (0.26)	0.46 (0.14)	0.030
Jan '96	2.81 (0.51)	1.84 (0.29)	0.001	0.73 (0.17)	0.27 (0.10)	0.001
Mar '96	4.01 (0.81)	1.83 (0.34)	0.001	0.60 (0.18)	0.17 (0.08)	0.001
May '96	3.86 (0.76)	2.68 (0.56)	0.125	0.72 (0.21)	0.27 (0.07)	0.095
July '96	8.33 (1.58)	5.46 (0.94)	0.037	2.35 (0.51)	0.60 (0.13)	0.035

[†] Indicates the only instance where root length density was not significantly greater in 0-15-cm strata.

Figure 1 Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina.

Figure 2 Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina

Figure 3 Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina

Figure 4 Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina

Figure 5 Fine Root Productivity and Dynamics on a Forested Floodplain in South Carolina