

Fall 11-12-2003

Evolution of the Polarization of the Optical Afterglow of the Gamma-ray Burst GRB 030329

Dieter H. Hartmann

Department of Physics and Astronomy, Clemson University, hdieter@clemson.edu

Jochen Greiner

Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse

Sylvio Klose

Thüringer Landessternwarte Tautenburg - Tautenburg, Germany

Klaus Reisch

Universitäts-Sternwarte Göttingen

Hans Martin Schmid

Institut für Astronomie

See next page for additional authors

Follow this and additional works at: https://tigerprints.clemson.edu/physastro_pubs

Recommended Citation

Please use publisher's recommended citation.

This Article is brought to you for free and open access by the Physics and Astronomy at TigerPrints. It has been accepted for inclusion in Publications by an authorized administrator of TigerPrints. For more information, please contact kokeefe@clemson.edu.

Authors

Dieter H. Hartmann, Jochen Greiner, Sylvio Klose, Klaus Reisch, Hans Martin Schmid, Re'em Sari, Chryssa Kouveliotou, Arne Rau, Eliana Palazzi, and Christian Straubmeier

Evolution of the polarization of the optical afterglow of the gamma-ray burst GRB 030329

Jochen Greiner^{*}, Sylvio Klose[†], Klaus Reinsch[‡], Hans Martin Schmid^ℓ, Re'em Sari[§], Dieter H. Hartmann[¶], Chryssa Kouveliotou[§], Arne Rau^{*}, Eliana Palazzi[◇], Christian Straubmeier^{**}, Bringfried Stecklum[†], Sergej Zharikov^{††}, Gagrik Tovmassian^{††}, Otto Bärbantner[‡], Christoph Ries[‡], Emmanuel Jehin^{¶¶}, Arne Henden^{§§}, Anlaug A. Kaas^{‡‡}, Tommy Grav^{§§}, Jens Hjorth^{ℓℓ}, Holger Pedersen^{ℓℓ}, Ralph A.M.J. Wijers[#], A. Kaufer^{¶¶}, Hye-Sook Park^{◇◇}, Grant Williams^{##}, Olaf Reimer^{##}

^{*}Max-Planck-Institut für extraterrestrische Physik, 85741 Garching, Germany

[†]Thüringer Landessternwarte, 07778 Tautenburg, Germany

[‡]Universitäts-Sternwarte Göttingen, 37083 Göttingen, Germany

^ℓInstitut für Astronomie, ETH Zürich, 8092 Zürich, Switzerland

[§]California Institute of Technology, Theoretical Astrophysics 130-33, Pasadena, CA 91125, USA

[¶]Clemson University, Department of Physics and Astronomy, Clemson, SC 29634, USA

^{§§}NSSTC, SD-50, 320 Sparkman Drive, Huntsville, AL 35805, USA

[◇]Istituto di Astrofisica Spaziale e Fisica Cosmica, CNR, Sezione di Bologna, 40129 Bologna, Italy

^{**}I. Physikalisches Institut, Universität Köln, 50937 Köln, Germany

^{††}Instituto de Astronomia, UNAM, 22860 Ensenada, Mexico

^{‡‡}Wendelstein-Observatorium, Universitätssternwarte, 81679 München, Germany

^{¶¶}European Southern Observatory, Alonso de Cordova 3107, Vitacura, Casilla 19001, Santiago 19, Chile

^{§§}Universities Space Research Association, U.S. Naval Observatory, P.O. Box 1149, Flagstaff, AZ 86002, USA

^{##}Nordic Optical Telescope, 38700 Santa Cruz de La Palma, Spain

^{§§}University of Oslo, Institute for Theoretical Astrophysics, 0315 Oslo, Norway, and Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

^{ℓℓ}Astronomical Observatory, NBIfAFG, University of Copenhagen, 2100 Copenhagen, Denmark

[#]Astronomical Institute Anton Pannekoek, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

^{◇◇}Lawrence Livermore National Laboratory, University of California, P.O. Box 808, Livermore, CA 94551, USA

^{##}MMT Observatory, University of Arizona, Tucson, AZ 85721, USA

^{##}Theoretische Weltraum-und Astrophysik, Ruhr-Universität Bochum, 44780 Bochum, Germany

The association of a supernova with GRB 030329^{1,2} strongly supports the collapsar model³ of γ -ray bursts (GRBs), where a relativistic jet⁴ forms after

the progenitor star collapses. Such jets cannot be spatially resolved because of their cosmological distances. Their existence is conjectured based on breaks in GRB afterglow light curves and the theoretical desire to reduce the GRB energy requirements. Temporal evolution of polarization^{5,6,7} may provide independent evidence for the jet structure of the relativistic outflow. Small-level polarization ($\sim 1\text{-}3\%$)⁸⁻¹⁷ has been reported for a few bursts, but the temporal evolution of polarization properties could not be established. Here, we report polarimetric observations of the afterglow of GRB 030329 with high signal-to-noise and high sampling frequency. We establish the polarization light curve, detect sustained polarization at the percent level, and find significant variability. The data imply that the afterglow magnetic field has small coherence length and is mostly random, probably generated by turbulence, in contrast with the high polarization detected in the prompt γ -rays from GRB 021206¹⁸. Our results suggest a different structure and origin of the magnetic field in the prompt vs. afterglow emission regions.

GRB 030329 triggered the High Energy Transient Explorer, HETE-II, on March 29, 2003 (11:37:14.67 UT)¹⁹. The discovery of the burst optical afterglow^{20,21} was quickly followed by a redshift measurement²² for the burster of $z=0.1685$ (~ 800 Mpc), thus making GRB 030329 the second closest long-duration²³ GRB ever studied, after GRB 980425³. The proximity of GRB 030329 resulted in very bright prompt and afterglow emission, leading to the best-sampled afterglow to date. Detailed optical spectroscopy revealed an underlying supernova (SN 2003dh)^{1,2} with an astonishing spectral similarity to SN 1998bw (at $z=0.0085$, associated with GRB 980425³), thus strongly supporting the link of long-duration GRBs with core-collapse supernovae.

The apparent brightness of GRB 030329 also afforded a unique opportunity to study the temporal evolution of polarization in the afterglow phase. Previous single measurements found low-level (1-3%) polarization⁸⁻¹⁵ in optical afterglows, and the only reports on variable polarization^{16,17} were based on few measurements with different instruments and modest signal-to-noise. We have overcome the previous sampling limitations with 31 polarimetric observations of the afterglow of GRB 030329 obtained with the same instrumentation (plus few more with different instruments) over a time period of 38 days for a total of an unprecedented ~ 50 hours of observations with an 8 m telescope. For the first time we establish a polarization light curve for an optical gamma-ray burst afterglow.

We performed relative photometry, and derived from each pair of simultaneous measurements at orthogonal angles the Stokes parameters U and Q. In order to obtain the intrinsic polarization of the GRB afterglow, we have to correct for Galactic interstellar polarization (mostly due to dust). We performed imaging polarimetry to derive the polarization parameters of seven stars in the field of GRB 030329, and obtained an interstellar (dust) polarization correction of 0.45% at position angle 155deg. Subtraction of the mean foreground polarization was performed in the Q/U plane ($Q_{fp}=0.0027\pm 0.0013$, $U_{fp}=-0.0033\pm 0.0017$). Figure 1 shows an R band image of the GRB 030329 field with the polarization "vectors" superimposed.

The temporal evolution of the degree and angle of polarization together with the R band photometry is shown in Figure 2. This figure demonstrates the presence of non-zero polar-

ization, $\Pi \sim 0.3\text{-}2.5\%$ throughout a 38-day period, with significant variability in degree and angle on time scales down to hours. Further, the spectropolarimetric data of the first three nights as well as the simultaneous R and K band imaging polarimetry during the second night show that the relative polarization and the position angle are wavelength independent (within the measurement errors of about 0.1%) over the entire spectral range. These data imply that polarization due to dust in the host galaxy of GRB 030329 does not exceed $\sim 0.3\%$. Dust destruction in the vicinity of the gamma-ray burster due to the strong radiation field would result in a monotonically decreasing polarization degree at constant position angle, which is not observed during the first days. We therefore conclude that the bulk of the observed variability is intrinsic to the afterglow.

Figure 2 shows that while the polarization properties show substantial variability (for which no simple empirical relationship is apparent), the R band flux is a sequence of power laws. During each of the power law decay phases the polarization is of order few percent, different from phase to phase, and variable within the phase, but not in tandem with the "bumps and wiggles" in the light curve. We observe a decreasing polarization degree shortly after the light curve break at ~ 0.4 days (as determined from optical^{21,24} and X-ray data^{25,26}). Rapid variations of polarization occur ~ 1.5 days after the burst, and could be related to the end of the transition period towards a new power law phase starting at ~ 1.7 days. Polarization eventually rises to a level of $\sim 2\%$, which remains roughly constant for another two weeks. At late time the underlying Type Ic SN 2003dh^{1,2} increasingly contributes to the total light and probably also to the observed polarization properties. Asymmetries in this type of supernova can produce polarization of order 1%^{27,28}, as we observe towards the end of our campaign.

GRB 030329 belongs to a growing group of bursts for which densely monitored afterglow light curves show significant "bumps" and "wiggles" relative to a simple power law decay (most notably GRB 021004 and GRB 011211). These bumps and wiggles complicate the interpretation of the polarization properties. While the rapid decrease in polarization degree during the first night is consistent with the model predictions^{5,6,7}, the position angle changes are not, and thus it remains to be proven whether or not these early polarization data support the break at 0.4 days as due to a jet. A connection to theoretical models throughout the entire period covered in Fig. 2 is a daunting task. A detailed comparison of our data with characteristic features of various theoretical models is beyond the scope of this paper, and will be presented elsewhere.

In summary, our data constitute the most complete and dense sampling of the polarization behaviour of a GRB afterglow to date. For GRB 030329 we conclude that the afterglow polarization probably did not rise above $\sim 2.5\%$ at any time, and that the polarization did not correlate with the flux. The low level of polarization implies that the components of the magnetic field parallel and perpendicular to the shock do not differ by more than $\sim 10\%$, and suggests an entangled magnetic field, probably amplified by turbulence behind shocks, rather than a pre-existing field. This is in contrast with the high level of polarization detected in the prompt γ -rays from GRB 021206¹⁸ and suggests a different structure and origin of the magnetic field in the prompt vs. afterglow emission regions. Evolving polarization properties provide a unique diagnostic tool for GRB studies, and the extremely complex light curve of the optical afterglow of GRB 030329 emphasises that measurements should be carried out with high sampling frequency.


1. Stanek, K.Z., et al. Spectroscopic discovery of the supernova 2003dh associated with GRB 030329. *Astrophys. J.* 591, L17-L20 (2003)
2. Hjorth, J., et al. A very energetic supernova associated with the γ -ray burst of 29 March 2003. *Nature* 423, 847-850 (2003)
3. Woosley, S.E., Eastman, R.G., Schmidt, B.P. Gamma-ray bursts and type Ic supernova SN 1998bw. *Astrophys. J.* 516, 788-796 (1999)
4. Mszros, P. Theories of gamma-ray bursts. *Ann. Rev. Astron. Astrophys.* 40, 137-169 (2002)
5. Sari, R. Linear polarization and proper motion in the afterglow of beamed gamma-ray bursts. *Astrophys. J.* 524, L43-L46 (1999)
6. Gruzinov, A. Strongly polarized optical afterglows of gamma-ray bursts. *Astrophys. J.* 525, L29-L31 (1999)
7. Ghisellini, G., Lazzati, D. Polarization light curves and position angle variation of beamed gamma-ray bursts. *Mon. Not. R. Astron. Soc.* 309, L7-L11 (1999)
8. Hjorth, J., et al. Polarimetric constraints on the optical afterglow emission from GRB 990123. *Science* 283, 2073-2075 (1999)
9. Wijers, R.A.M.J., et al. Detection of polarization in the afterglow of GRB 990510 with the ESO Very Large Telescope. *Astrophys. J.* 523, L33-L36 (1999)
10. Covino, S., et al. GRB 990510: linearly polarized radiation from a fireball. *Astron. Astrophys.* 348, L1-L4 (1999)
11. Rol, E., et al. GRB 990712: First indication of polarization variability in a gamma-ray burst afterglow. *Astrophys. J.* 544, 707-711 (2000)
12. Björnsson, G., Hjorth, J., Pedersen, K., Fynbo, J.U. The afterglow of GRB 010222: A case of continuous energy injection. *Astrophys. J.* 579, L59-L62 (2002)
13. Masetti, N., et al. Optical and near-infrared observations of the GRB 020405 afterglow. *Astron. Astrophys.* 404, 465-481 (2003)
14. Covino, S., et al. Polarization evolution of the GRB 020405 afterglow. *Astron. Astrophys.* 400, L9-L12 (2003)
15. Bersier, D., et al. The strongly polarized afterglow of GRB 020405. *Astrophys. J.* 583, L63-L66 (2003)
16. Barth, A., et al. Optical spectropolarimetry of the GRB 020813. *Astrophys. J.* 584, L47-L51 (2003)

17. Rol, E., et al. Variable polarization in the optical afterglow of GRB 021004. *Astron. Astrophys.* 405, L23-L27 (2003)
18. Coburn, W., and Boggs, S.E. Polarization of the prompt (γ -ray emission from the (γ -ray burst of 6 December 2002. *Nature* 423, 415-417 (2003)
19. Vanderspek, R., Crew, G., Doty, J., Villasenor, J., Monelly, G. GRB 030329 (=H2652): A long, extremely bright GRB localized by HETE WXM and SCX. *GCN Circ.* 1997 (2003)
20. Price, P.A., et al. The bright optical afterglow of the nearby (γ -ray burst of 29 March 2003. *Nature* 423, 844-847 (2003)
21. Uemura, M., et al. Structure in the early afterglow light curve of the (γ -ray burst of 29 March 2003. *Nature* 423, 843-844 (2003)
22. Greiner, J., et al. Redshift of GRB 030329. *GCN Circ.* 2020 (2003)
23. Kouveliotou, C., et al. Identification of two classes of gamma-ray bursts. *Astrophys. J.* 413, L101-L104 (1993)
24. Burenin, R., et al. First hours of the GRB 030329 optical afterglow. *Astron. Lett.* 29, 9, 1-6 (2003)
25. Marshall, F.E., Markwardt, C., Swank, J.H. GRB 030329: fading X-ray afterglow with RXTE. *GCN Circ.* 2052 (2003)
26. Tiengo, A., et al. The X-ray afterglow of GRB 030329. *Astron. Astrophys.*, in press; astro-ph/0305564 (2003)
27. Wang, L., Howell, D. A., Hflich, P., and Wheeler, J. C. Bipolar supernova explosions. *Astrophys. J.* 550, 1030-1035 (2001)
28. Leonard, D. C., Filippenko, A. V., Chornock, R., and Foley, R. J. Photospheric-phase spectropolarimetry and nebular-phase spectroscopy of the peculiar type Ic supernova 2002ap. *Publ. Astron. Soc. Pac.* 114, 1333-1348 (2002)

Acknowledgements

This work is primarily based on observations collected at ESO, Chile, with additional data obtained at the German-Spanish Astronomical Centre Calar Alto, operated by the Max-Planck-Institute for Astronomy, Heidelberg, jointly with the Spanish National Commission for Astronomy, the NOT on La Palma, Canary Islands, and the Observatorio Astronomico National, San Pedro, Mexico. We are grateful to the staff at the Paranal, Calar Alto and NOT observatories, in particular A. Aguirre, M. Alises, S. Hubrig, A.O. Jaunsen, C. Ledoux, S. Pedraz, T. Szeifert, L. Vanzì and P. Vreeswijk for obtaining the service mode data reported here.

Figure 1: R band image of the field centred on the optical afterglow of GRB 030329. The 27 FORS1 (Focal Reducer and Low Dispersion Spectrograph at VLT/Antu) measurements are shown as "vectors" where the length is a measure of the polarization degree, and the orientation indicates the position angle. While the afterglow varies in degree as well as angle, the polarization of seven field stars is constant within 0.1% in polarization degree and 1.5 degree in position angle throughout the 38 days. Linear polarization was measured from sets of exposures with different retarder plate position angles. Imaging polarimetry was obtained during the first four nights (when the afterglow was brighter than 17th mag) from sixteen different retarder plate angles, and from eight angles thereafter. Since the FORS1 polarization optics allows determination of the degree of polarization to an accuracy of $\pm 3 \times 10^{-4}$ and of the polarization angle to ~ 0.2 deg, we consider the above variance of the field stars to represent the systematic error over the 38-day time period. Observations of polarimetric standard stars reproduced their tabulated values within 5%. The FORS1 retarder plate zero point angle of -1.2 degrees was subtracted from the polarization angle. The position angle has a systematic uncertainty of 1.5 deg, which was added in quadrature to the statistical errors.






Figure 2: Evolution of the polarization during the first 38 days. Top and middle panels show the polarization degree in percent and the position angle in degrees. The red data points are from imaging polarimetry with FORS1/VLT. Spectropolarimetry (blue symbols, 700-800 nm range) was performed during the first three nights. The green points were obtained with CAFOS at the 2.2m telescope at Calar Alto on March 30/31. The magenta data point was obtained with AFOSC at the 2.56m NOT telescope on 2003 April 2. The bottom panel shows the residual R band light curve after subtraction of the contribution of a power law $t^{-1.64}$ describing the undisturbed decay during the time interval 0.5-1.2 days after the GRB (i.e., after the early break at 0.4 days), thus leading to a horizontal curve. The symbols correspond to data obtained through: the literature (black), the 1m USNO telescope at Flagstaff (blue), the OAN Mexico (cyan), and FORS1/VLT (red). Lines indicate phases of power law decay, with the first one from early data21 (not shown). Yellow bars mark re-brightening transitions. Contributions from an underlying supernova (solid curved line) do not become significant until ~ 10 days after the GRB.