

8-1-2014

Low Vitamin D Status and Inadequate Nutrient Intakes of Elementary School Children in a Highly Educated Pacific Northwest Community

Simone Frei

Oregon State University, simone.frei@oregonstate.edu

Balz Frei

Oregon State University, balz.frei@oregonstate.edu

Gerd Bobe

Oregon State University, gerd.bobe@oregonstate.edu



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/).

Recommended Citation

Frei, S., Frei, B., & Bobe, G. (2014). Low Vitamin D Status and Inadequate Nutrient Intakes of Elementary School Children in a Highly Educated Pacific Northwest Community. *The Journal of Extension*, 52(4), Article 10. <https://tigerprints.clemson.edu/joe/vol52/iss4/10>

This Research in Brief is brought to you for free and open access by the Conferences at TigerPrints. It has been accepted for inclusion in The Journal of Extension by an authorized editor of TigerPrints. For more information, please contact kokeefe@clemson.edu.

Low Vitamin D Status and Inadequate Nutrient Intakes of Elementary School Children in a Highly Educated Pacific Northwest Community

Abstract

Are Extension healthy youth programs needed in highly educated U.S. communities? To answer this question, 175 children from four public elementary schools in Corvallis, Oregon, self-reported in a cross-sectional study their dietary intake, and 71 children provided a blood sample for measuring vitamin D concentrations. Most children had insufficient blood vitamin D and reported a diet insufficient in fiber, essential fatty acids, potassium, and vitamin E and excessive in saturated fatty acids and sodium. Our data indicate a need for Extension to provide youth programs in highly educated U.S. communities to improve nutrient intakes and vitamin D status in children.

Simone Frei
Healthy Youth
Program Manager
simone.frei@oregonstate.edu

Balz Frei
Distinguished
Professor
balz.frei@oregonstate.edu

Gerd Bobe
Assistant Professor
gerd.bobe@oregonstate.edu

Oregon State
University
Linus Pauling
Institute
Corvallis, Oregon

Introduction

The foundation for a healthy life is laid in childhood, as the body has increased demands of nutrients for growth and development of organs to achieve optimum function. Nutrient recommendations and guidelines have been established by experts and government officials for specific age groups, including 4-8 and 9-13 year-old children, stratified by sex (FNB & IOM, 2005; USDA & HHS, 2010). Vitamin D status depends mainly on sun exposure of the skin (Adams, Clemens, Parrish, & Holick, 1982); however, in areas with low sun exposure, such as the Pacific Northwest, dietary intake becomes very important. Various healthy youth programs have been provided by Extension professionals to improve nutrient intakes in U.S. elementary school-aged children with a focus on low-income families because of their limited resources; however, improvements were limited (Rabe, Ohri-Vachaspati, & Scheer, 2006; Roth-Yousey, Caskey, May, & Reicks, 2007; Jensen, Kattelmann, Ren, & Wey, 2009).

In U.S. population surveys, most elementary school-aged children report nutrient intakes low in fiber and vitamin E and high in saturated fat and sodium (Rhodes, Clemens, Goldman, Lacombe, & Moshfegh, 2012) with insufficient vitamin D concentrations in blood (Kumar, Muntner, Kaskel, Hailpern, & Melamed, 2009; Mansbach, Ginde, & Cargo, 2009), regardless of household income. These studies, however, do not break down nutrient intakes and vitamin D concentrations by communities

but rather by demographics of individuals. Some of the challenges to achieve adequate nutrient intakes and vitamin D status in school-aged children are: a) limited access to nutrient-rich foods and beverages in the community and at home (Ogden, Lamb, Carroll, & Flegal, 2010; SEARCH, 2007); b) school meals that are low in whole grains, legumes, fiber, and vitamin E and high in saturated fat and sodium (Condon, Crepinsek, & Fox, 2009; Gordon & Fox, 2007, 2012); c) limited sun exposure (Bose et al., 2013), and d) the consumption of high-simple carbohydrate, low-fiber, and low-vitamin-D foods and beverages, which increases as children become teenagers (Ballew, Kuester, & Gillespie, 2000; Bowman, Gortmaker, Ebbelin, Pereira, & Ludwig, 2004; Evans, Springer, Evans, Ranjit, & Hoelscher, 2010).

Most public funding and scientific studies for Extension programs focus on communities with limited access to nutrient-rich foods and beverages in the community and at home and lower education level because of their limited resources. Given budget constraints, the question arises: Are Extension healthy youth programs needed in highly educated U.S. communities? To answer, we need to know whether vitamin D status and nutrient intakes are adequate in highly educated communities. Such communities have adequate access to nutrient-rich foods and beverages, because multiple grocery stores with a wide variety of fruits and vegetables are in close proximity to schools and homes. Information about the vitamin D status and nutrient intake of children in such communities is lacking. Therefore, we compared in a cross-sectional study food and nutrient intake of elementary school-aged children from four public elementary schools in Corvallis, a college town in the Pacific Northwest, with dietary guidelines and their blood vitamin D levels. Using generally used vitamin D cut-off values for children (Gilbert-Diamond et al., 2010), the proportion of children with vitamin D levels (sum of 25-hydroxyvitamin D2 and 25-hydroxyvitamin D3) below 30 and 20 ng/mL were considered insufficient and deficient, respectively.

Methods

The study was reviewed and approved by the Oregon State University Institutional Review Board (IRB Permit Number: 4841; "Micronutrient Intake of Elementary School Children"). Corvallis (population size: 52,396 in 2010; 83% White, 8% Asian, 1% Black or African American, and 8% Hispanics or other), Oregon, is a town with one of the highest education levels in the U.S.: 30.2% of the population have completed a bachelor's degree (U.S. average: 17.5%), and 26.4% have completed a graduate degree (U.S. average: 9.8%) (Corvallis Oregon Demographics, 2013). In 2010, the median household income in Corvallis was \$47,339, which was below the median U.S. household income of \$55,970.

After receiving approval from the school district and school principals, four public elementary schools in Corvallis participated in the study. Grocery stores that carry a broad variety of fruits and vegetables are within 2 miles of each of the four elementary schools. The proportion of children from low income households, as defined by the Oregon Department of Education, was 11%, 17%, 19%, and 48% for the four schools, respectively (ODE, 2013). All children who attended the schools, aged 5-11 years, were eligible.

The study staff (the program manager and a dietetics student) visited each class room in the four elementary schools, explained the study objective and demonstrated to children how to fill out the Block Kids Food Screener (BKFS) and how parents would collect blood by finger prick and administer it

to the vitamin D test. The visit of the first school took place in October 2009 and did not include the vitamin D test. The remaining three schools were visited between April and December 2011 and included the vitamin D test. At the end of the demonstration, students received an information package for their parents that included information on the study, how to fill out the BKFS, how to administer the vitamin D test, as well as informed parent consent and student assent forms.

Upon return of the signed forms, the vitamin D test was sent out by ZRT Laboratory (Beaverton, OR) to parents, who completed the test with their child and sent it back to ZRT Laboratory, a certified, diagnostic laboratory for measuring human blood samples using standard reference materials. Dried blood spot samples for the vitamin D test were returned by 71 children (7% of children in the three elementary schools), and the results were included in the statistical analysis. Of the 71 children with a vitamin D blood test, 38 were 5-8 years old (13 boys and 25 girls), and 33 were 9-11 years old (14 boys and 19 girls). Nearly all children (67 of 71 children) with a blood vitamin D test had at the same time also filled out the BKFS. Vitamin D concentrations were measured as previously described (Newman et al., 2009). The inter- and intra-assay coefficients of variation were 10.4% and 8.7%, respectively.

The BKFS, developed by Nutrition Quest (Berkeley, California), is a 41-item, two-page food frequency questionnaire (FFQ) that asks for frequency (six categories from none to every day) and quantity of dietary intakes (four categories) of the most commonly consumed food items by children during the past week (Hunsberger, O'Malley, Block, & Norris, 2012). The food screener has been described and validated for 2-17 year-old U.S. children (Garcia-Dominic et al., 2012; Hunsberger et al., 2012). We did not specify that the children had to fill out the BKFS without the help of adults. A total of 179 children, which accounted for 13% of all children in the four elementary schools, completed the BKFS and returned it by mail to the study staff.

The completed BKFS were analyzed by Nutrition Quest for food servings and nutrient intakes. Food codes were linked to the USDA Food and Nutrient Database for Dietary Studies and the USDA My Pyramid Equivalents Data Base to calculate food group and nutrient intakes. Results from four BKFS were not included in the statistical analysis because the reported energy consumption was more than three standard deviations different from the mean. Of the 175 children included in the analysis, 75 were 5-8 years old (28 boys and 47 girls), and 100 were 9-11 years old (45 boys and 55 girls).

Because FFQs are known to underestimate food and nutrient intake, the calculated food group and nutrient intakes were calorie-adjusted and expressed per 1,000 Kcal. To calculate the proportion of children not meeting recommendations, calorie-adjusted intakes were multiplied by the estimated energy requirements for each age and gender group (FNB & IOM, 2005) and then compared to governmental dietary guidelines and recommendations (USDA & HHS, 2010). The recommended food group intakes differ by caloric consumption, and the nutrient intakes differ by sex and age group and, thus, are not shown.

Statistical analyses were performed using SAS version 9.2 software (SAS Institute, 2009). Food group servings, nutrient intakes, and vitamin D concentrations were compared between age groups (5-8 year-old *versus* 9-11 year-old) using PROC GLM for calorie-adjusted values and PROC GLIMMIX for proportions meeting recommendations. Values presented in the manuscript are raw means and their

standard errors (SEM). We tested the effects of season and school on nutrient intakes and vitamin D status. The effects were not significant and were not included in the model. All statistical tests were two-sided. Statistical significance was declared at $P = 0.05$.

Results and Discussion

The results of our cross-sectional study document that low vitamin D status and inadequate nutrient intake in elementary school-aged children is prevalent even in a highly educated community with adequate access to stores with nutrient-rich foods and beverages. Regarding vitamin D status (Table 1), $69 \pm 6\%$ of children had vitamin D concentrations < 30 ng/mL, which are considered insufficient, and $8 \pm 3\%$ had vitamin D concentrations considered deficient (< 20 ng/mL). Older children had lower vitamin D concentrations than younger children. To explore the relationship between blood vitamin D concentrations and food group intakes, we used multivariate regression. We did not have dietary vitamin D results. Consumption of dairy products but no other food group was significantly associated with increased serum vitamin D concentrations ($\beta = 2.2 \pm 1.0$ cups/1,000 Kcal; $P = 0.04$). Simulation studies indicate that consumption of dairy products is important for providing sufficient dietary vitamin D, calcium, phosphorus, magnesium, and potassium to U.S. children (Gao, Wilde, Lichtenstein, & Tucker, 2006; Fulgoni, Keast, Auestad, & Quann, 2011).

Table 1.

Blood Vitamin D Concentrations of Elementary School Children in a Highly Educated Community in the Pacific Northwest

Serum Vitamin D	5-8 years old (n = 38)	9-11 years old (n = 33)	P-value ¹
	Mean \pm SEM		
Concentrations (ng/mL)	28.0 \pm 1.0	24.7 \pm 1.0	0.03
Range (ng/mL)	15 – 42	12 - 37	
25(OH) Vitamin-D Levels	% of Children \pm SEM		
< 15 ng/mL	0 \pm 0	6 \pm 4	0.21
< 20 ng/mL	5 \pm 4	12 \pm 6	0.32
< 30 ng/mL	61 \pm 8	79 \pm 7	0.11
1 P-value for difference between age groups			

Older children also consumed fewer dairy products compared to younger children, and $26 \pm 3\%$ of all children did not meet the recommendations (adjusted for total caloric intake) for the consumption of dairy products (USDA & HHS, 2010; Table 2). Coinciding with the lower consumption of dairy products in older children, $45 \pm 5\%$ of the older children consumed calcium below dietary guidelines, and $12 \pm 3\%$ of older children consumed less phosphorus than recommended (Table 3).

Table 2.

Self-Reported Daily Food Group Consumption of Elementary School Children in a Highly Educated Community in the Pacific Northwest

Food Groups	5-8 years old (n = 75)	9-11 years old (n = 100)	P-value¹
Consumption/Content	Mean ± SEM²		
Fruit (cup/1,000 Kcal))	1.39 ± 0.08	1.49 ± 0.07	0.30
Vegetables (cup/1,000 Kcal)	0.87 ± 0.04	1.13 ± 0.05	0.0005
Potatoes (cup/1,000 Kcal)	0.158 ± 0.013	0.220 ± 0.014	0.003
Legumes (cup/1,000 Kcal)	0.074 ± 0.011	0.088 ± 0.010	0.39
Other Vegetables (cup/1,000 Kcal)	0.64 ± 0.03	0.82 ± 0.04	0.002
Whole Grains (oz/1,000 Kcal)	0.68 ± 0.04	0.61 ± 0.04	0.26
Meat/Poultry/Fish (oz/1,000 Kcal)	1.69 ± 0.09	1.67 ± 0.08	0.89
Dairy (cup/1,000 Kcal)	1.75 ± 0.08	1.55 ± 0.07	0.05
Not Meeting Recommendations	% of Children ± SEM³		
Fruit	8 ± 3	8 ± 3	1
Vegetables	33 ± 5	12 ± 3	0.001
Potatoes	69 ± 5	39 ± 5	0.0001
Legumes	100 ± 0	100 ± 0	1
Other Vegetables	23 ± 5	13 ± 3	0.10
Whole Grains	83 ± 4	74 ± 4	0.18
Meat/Poultry/Fish	23 ± 5	19 ± 4	0.55
Dairy	24 ± 5	27 ± 4	0.65

¹ P-value for difference between age groups

² The estimated food group intake was calculated by multiplying the self-reported consumption (in cup or oz/1,000 Kcal) by the estimated energy requirement.

³ % of children was calculated by comparing the recommended with the

estimated food group consumption per 1,000 Kcal for the respective age and gender group.

Table 3.

Self-Reported Daily Mineral Consumption of Elementary School Children in a Highly Educated Community in the Pacific Northwest

Minerals	5-8 years old 6- (n = 75)	9-11 years old 10- (n = 100)	P-value¹
Consumption/Content	Mean ± SEM²		
Calcium (mg/1,000 Kcal)	704 ± 21	665 ± 18	0.17
Phosphorus (mg/1,000 Kcal)	795 ± 14	762 ± 13	0.08
Magnesium (mg/1,000 Kcal)	152.7 ± 2.1	156.1 ± 2.3	0.29
Iron (mg/1,000 Kcal)	7.95 ± 0.22	8.22 ± 0.23	0.40
Zinc (mg/1,000 Kcal)	6.73 ± 0.13	6.65 ± 0.12	0.62
Copper (µg/1,000 Kcal)	561 ± 10	602 ± 9	0.002
Selenium (µg/1,000 Kcal)	54.6 ± 0.9	50.2 ± 0.8	0.0002
Potassium (mg/1,000 Kcal)	1651 ± 28	1742 ± 32	0.04
Sodium (mg/1,000 Kcal)	1524 ± 25	1509 ± 22	0.66
Not Meeting Guidelines	% of Children ± SEM³		
Calcium (<EAR)	16 ± 4	45 ± 5	0.0001
Phosphorus (<EAR)	0 ± 0	12 ± 3	0.001
Magnesium (<EAR)	0 ± 0	3 ± 2	0.26
Iron (<EAR)	0 ± 0	0 ± 0	1
Zinc (<EAR)	0 ± 0	0 ± 0	1
Copper (<EAR)	0 ± 0	0 ± 0	1
Selenium (<EAR)	0 ± 0	0 ± 0	1
Potassium (<AI)	100 ± 0	99 ± 1	1
Sodium (>UL)	92 ± 3	87 ± 3	0.30

EAR: estimated average requirement; AI: adequate intake; UL: tolerable upper intake level

¹ P-value for difference between age groups

² The estimated mineral intake was calculated by multiplying the self-reported

consumption (in mg or $\mu\text{g}/1,000$ Kcal) by the estimated energy requirement. 3 % of children was calculated by comparing the recommended with the estimated mineral consumption per for the respective age and gender group.

Besides vitamin D, all children and $78 \pm 3\%$ of children did not meet the recommendations for legume and whole grain consumption, respectively (Table 2). In response, $70 \pm 3\%$ of children consumed less fiber than recommended (Table 4). Besides fiber, legume consumption is associated with increased intake of potassium, phosphorus, magnesium, and folate (Papanikolaou, & Fulgoni, 2008). Nearly all children ($99 \pm 0.6\%$ and $89 \pm 2\%$, respectively) consumed less potassium and more sodium than recommended (Table 3). Targeting legume and whole-grain consumption together may improve nutrient intakes of children.

Table 4.

Self-Reported Daily Macronutrient Consumption of Elementary School Children in a Highly Educated Community in the Pacific Northwest

Macronutrients	5-8 years old (n = 75)	9-11 years old (n = 100)	P-value ¹
Energy (kcal)	Mean \pm SEM²		
Reported Intake	1,074 \pm 27	1,227 \pm 39	0.003
Consumption/Content (g/1,000 Kcal)			
Proteins	43.7 \pm 0.7	42.5 \pm 0.6	0.19
Carbohydrates	123.4 \pm 1.6	129.0 \pm 1.5	0.01
Sugar	81.8 \pm 1.9	87.5 \pm 2.0	0.05
Sugar in Foods/Beverages	64.7 \pm 1.6	69.5 \pm 1.6	0.03
Added Sugar	17.1 \pm 0.7	17.9 \pm 1.0	0.51
Sugar from Sodas	1.58 \pm 0.44	3.80 \pm 0.66	0.01
Starch	30.7 \pm 1.1	30.3 \pm 1.3	0.83
Fiber	10.96 \pm 0.30	11.24 \pm 0.28	0.48
Total Fat	36.8 \pm 0.6	34.9 \pm 0.5	0.02
Saturated Fat	13.43 \pm 0.29	12.77 \pm 0.23	0.07
Linoleic Acid (mg/1,000 Kcal)	4.78 \pm 0.08	4.71 \pm 0.09	0.58
Linolenic Acid (mg/1,000 Kcal)	0.527 \pm 0.010	0.522 \pm 0.009	0.74

Cholesterol (mg/1,000 Kcal)	137.9 ± 5.1	120.8 ± 3.9	0.008
Consumption (% of calories)			
Total Fat (% of calories)	33.2 ± 0.5	31.4 ± 0.5	0.02
Saturated Fat (% of calories)	12.09 ± 0.26	11.49 ± 0.21	0.07
Not Meeting Recommendations	% of Children ± SEM³		
Protein (<RDA)	0 ± 0	0 ± 0	1
Carbohydrate (<RDA)	0 ± 0	0 ± 0	1
Fiber (<IOM)	60 ± 6	78 ± 4	0.01
Total Fat (>AMDR)	35 ± 6	20 ± 4	0.03
Saturated Fat (>DG)	83 ± 4	80 ± 4	0.66
Linoleic Acid (<AI)	99 ± 1	93 ± 3	0.11
Linolenic Acid (<AI)	71 ± 5	84 ± 4	0.04
Cholesterol (>DG)	13 ± 4	8 ± 3	0.26
<p>RDA: recommended dietary allowance; IOM: Institute of Medicine; AMDR: acceptable macronutrient distribution range; DG: dietary guidelines; AI: adequate intake</p> <p>1 P-value for difference between age groups</p> <p>2 The estimated macronutrient intake was calculated by multiplying the self-reported consumption (in g or mg/1,000 Kcal or % calories) by the estimated energy requirement.</p> <p>3 % of children was calculated by comparing the recommended with the estimated macronutrient consumption per 1,000 Kcal or % calories for the respective age and gender group.</p>			

We also observed inadequate intakes of lipids and fat soluble vitamins. Only one child met the intake guidelines for vitamin E, and 22 ± 3% of children consumed less vitamin K than recommended (Table 5). Furthermore, 95 ± 2% of children did not meet the guidelines for linoleic acid and 78 ± 3% for linolenic acid (Table 4). Conversely, 81 ± 3% of children exceeded the guidelines of getting less than 10% of calories from saturated fat, and 26 ± 3% and 10 ± 2% of children, respectively, consumed more total fat (>35% of calories) and cholesterol (>300 g/day) than recommended (Table 4).

Table 5.

Self-Reported Daily Vitamin Consumption of Elementary School Children in a Highly Educated Community in the Pacific Northwest

	5-8 years old	9-11 years old	
--	----------------------	-----------------------	--

Vitamins	(n = 75)	(n = 100)	P-value¹
Consumption/Content	Mean ± SEM²		
Vitamin A (µg RAE/1,000 Kcal)	494 ± 16	499 ± 14	0.82
Vitamin C (mg/1,000 Kcal)	62.8 ± 2.8	76.6 ± 2.9	0.001
Vitamin E (mg/1,000 Kcal)	2.76 ± 0.05	2.96 ± 0.05	0.005
Thiamin (mg/1,000 Kcal)	0.877 ± 0.016	0.866 ± 0.015	0.60
Riboflavin (mg/1,000 Kcal)	1.40 ± 0.03	1.32 ± 0.03	0.08
Niacin (mg/1,000 Kcal)	10.19 ± 0.22	10.2 ± 0.21	0.94
Vitamin B6 (mg/1,000 Kcal)	1.084 ± 0.022	1.106 ± 0.020	0.46
Folate (µg DFE/1,000 Kcal)	320 ± 11	314 ± 10	0.73
Vitamin B12 (µg/1,000 Kcal)	3.37 ± 0.11	3.11 ± 0.09	0.06
Vitamin K (µg/1,000 Kcal)	60.6 ± 4.7	77.7 ± 5.7	0.03
Not Meeting Guidelines	% of Children ± SEM³		
Vitamin A (RAE; <EAR)	0 ± 0	1 ± 1	1
Vitamin C (<EAR)	0 ± 0	0 ± 0	1
Vitamin E (<EAR)	99 ± 1	100 ± 0	0.43
Thiamin (<EAR)	0 ± 0	0 ± 0	1
Riboflavin (<EAR)	0 ± 0	0 ± 0	1
Niacin (<EAR)	0 ± 0	0 ± 0	1
Vitamin B6 (<EAR)	0 ± 0	0 ± 0	1
Folate (DFE; <EAR)	0 ± 0	0 ± 0	1
Vitamin B12 (<EAR)	0 ± 0	0 ± 0	1
Vitamin K (<AI)	32 ± 5	15 ± 4	0.01
<p>RAE: retinol activity equivalents; DFE: dietary folate equivalents; EAR: estimated average requirement; AI: adequate intake</p> <p>¹ P-value for difference between age groups</p> <p>² The estimated vitamin intake was calculated by multiplying the self-reported consumption (in mg or µg/1,000 Kcal) by the estimated energy requirement.</p> <p>³ % of children was calculated by comparing the recommended with the estimated vitamin consumption per for the respective age and gender group.</p>			

Implications for Extension

Adequate nutrient status in childhood is critical for optimum development and function of mind and body throughout life. Our Extension project data indicate that, even in a highly educated community, low vitamin D status and inadequate nutrient intake (insufficient in fiber, essential fatty acids, potassium, and vitamin E while excessive in saturated fatty acids and sodium) are prevalent in elementary school children and worse in older children. Therefore, nutrition programs should be inclusive for all youth populations, not just the low-income and other under-served groups that we are often encouraged to focus on. Based on our results and input from elementary school-aged children and their parents, we developed inclusive Extension educational programs with hands-on activities for the youth and their families. These include family and youth gardening and cooking programs, grocery store tours for healthful meals on a budget, kid-friendly cookbooks, and healthful recipes for food that is commonly available in food banks. These programs are currently under evaluation.

Acknowledgements

The authors thank Nova Elwood for her assistance in recruiting children, study management, and data collection and Thushanthi Perera for her assistance in manuscript preparation. We would like to thank the principals and staff of Adams Elementary School, Franklin Elementary School, Hoover Elementary School, and Wilson Elementary School (all in Corvallis, Oregon) for allowing us to recruit children from their school. The study was supported financially by USANA Health Sciences, Inc. (Salt Lake City, Utah).

References

- Adams, J. S., Clemens, T. L., Parrish, J. A., & Holick, M. F. (1982). Vitamin-D synthesis and metabolism after ultraviolet irradiation of normal and vitamin-D-deficient subjects. *New England Journal of Medicine*, 306(12), 722-735.
- Ballew, C., Kuester, S., & Gillespie, C. (2000). Beverage choices affect adequacy of children's nutrient intakes. *Archives of Pediatrics & Adolescent Medicine*, 154(11), 1148-1152. Retrieved from: <http://archpedi.jamanetwork.com/article.aspx?articleid=352051>
- Bose, S., Breyse, P. N., McCormack, M.C., Hansel, N. N., Rusher, R. R., Matsui, E., Peng, R., Curtin-Brosnan, J., & Diette, G. B. (2013). Outdoor exposure and vitamin D levels in urban children with asthma. *Nutrition Journal*, 12(81), 1-7. Retrieved from: <http://www.nutritionj.com/content/pdf/1475-2891-12-81.pdf>
- Bowman, S. A., Gortmaker, S. L., Ebbelin, C. B., Pereira, M. A., & Ludwig, D. S. (2004). Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics*, 113(1), 112-118. Retrieved from: <http://pediatrics.aappublications.org/content/113/1/112.full.pdf+html>
- Condon, E. M., Crepinsek, M. K., & Fox, M. K. (2009). School meals: types of foods offered to and consumed by children at lunch and breakfast. *Journal of the American Dietetics Association*, 109(2), S67-S78.

Corvallis Oregon Demographics (2013). *Corvallis Demographics Summary*. Retrieved from:
<http://www.clrsearch.com/Corvallis-Demographics/OR/>

Evans, A. E., Springer, A. E., Evans, M. H., Ranjit, N., & Hoelscher, D. M. (2010). A descriptive study of beverage consumption among an ethnically diverse sample of public school students in Texas. *Journal of the American College of Nutrition*, 29(4), 387-396. Retrieved from:
<http://intl.jacn.org/cgi/reprint/29/4/387.pdf>

Food and Nutrition Board (FNB), & Institute of Medicine (IOM) of the National Academies (2005). *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. The National Academy Press: Washington, DC.

Fulgoni, V. L. 3rd, Keast, D. R., Auestad, N., & Quann, E. E. (2011). Nutrients from dairy foods are difficult to replace in diets of Americans: food pattern modeling and an analyses of the National Health and Nutrition Examination Survey 2003-2006. *Nutrition Research* 31(10), 759-765. Retrieved from:
<http://www.sciencedirect.com/science/article/pii/S0271531711001886#>

Gao, X., Wilde, P. E., Lichtenstein, A. H., & Tucker, K.L. (2006). Meeting adequate intake for dietary calcium without dairy foods in adolescents aged 9 to 18 years (National Health and Nutrition Examination Survey 2001-2002). *Journal of the American Dietetics Association*, 106(11), 1759-1765. Retrieved from: <http://www.sciencedirect.com/science/article/pii/S0002822306018438#>.

Garcia-Dominic, O., Treviño, R. P., Echon, R. M., Mobley, C., Block, T., Bizzari, A., & Michalek, J. (2012). Improving quality of food frequency questionnaire response in low-income Mexican American children. *Health Promotion Practice*, 13(6), 763-771.

Gilbert-Diamond, D., Baylin, A., Mora-Plazas, M., Marin, C., Arsenault, J. E., Hughes, M. D., Willet, W. C., & Villamor, E. (2010). Vitamin D deficiency and anthropometric indicators of adiposity in school-age children: a prospective study. *American Journal of Clinical Nutrition*, 92(6), 1446-1451. Retrieved from: <http://ajcn.nutrition.org/content/92/6/1446.full.pdf+html>

Gordon, A., & Fox, M. K. (2007). School Nutrition Dietary Assessment Study – III. *Summary of findings*. United States Department of Agriculture. Food and Nutrition Service. Office of Research and Analysis. November 2007. Retrieved from:
<http://www.fns.usda.gov/Ora/menu/Published/CNP/Files/SNDIII-SummaryofFindings.pdf>

Gordon, A., & Fox, M. K. (2012). School Nutrition Dietary Assessment Study – IV. *Summary of findings*. United States Department of Agriculture. Food and Nutrition Service. Office of Research and Analysis. November 2012. Retrieved from: http://www.mathematica-mpr.com/publications/PDFs/nutrition/snda-iv_findings.pdf

Hunsberger, M., O'Malley, J., Block, T., & Norris, J. C. (2012). Relative validation of Block Kids Food Screener for dietary assessment in children and adolescents. *Maternal & Child Nutrition*, 2012; Sep 24. doi: 10.1111/j.1740-8709.2012.00446.x. [Epub ahead of print].

Jensen, B., Kattelman, K., Ren, C., & Wey, H. (2009). The efficacy of KidQuest: A nutrition and physical activity curriculum for 5th and 6th grade youth. *Journal of Extension* [On-line], 47(3) Article 3FEA4. Available at: <http://www.joe.org/joe/2009june/a4.php>

- Kumar, J., Muntner, P., Kaskel, F. J., Hailpern, S. M., & Melamed, M. L. (2009). Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001-2004. *Pediatrics*, 124(3), e362-e370. Retrieved from: <http://pediatrics.aappublications.org/content/124/3/e362.full.pdf+html>
- Mansbach, J. M., Ginde, A. A., & Cargo, C. A. Jr. (2009). Serum 25-hydroxyvitamin D levels among US children aged 1 to 11 years: do children need more vitamin D? *Pediatrics*, 124(5), 1404-1410. Retrieved from: <http://pediatrics.aappublications.org/content/124/5/1404.full.pdf+html>
- Newman, M. S., Brandon, T. R., Groves, M. N., Gregory, W. L., Kapur, S., & Zava, D. T. (2009). A liquid chromatography/tandem mass spectrometry method for determination of 25-hydroxy vitamin D2 and 25-hydroxy vitamin D3 in dried blood spots: a potential adjunct to diabetes and cardiometabolic risk screening. *Journal of Diabetes Science and Technology*, 3(1), 156-162. Retrieved from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2769851/>
- Ogden, C. L., Lamb, M. M., Carroll, M. D., & Flegal, K. M. (2010). *Obesity and socioeconomic status in children and adolescents: United States 2005-2008*. NCHS data brief no 51, MD: National Center for Health Statistics December 2010. Retrieved from: <http://www.cdc.gov/nchs/databriefs/db51.pdf>
- Oregon Department of Education (ODE) (2013). *Report cards*. Retrieved from: <http://www.ode.state.or.us/data/reportcard/reports.aspx>
- Papanikolaou, Y., & Fulgoni, V. L. 3rd (2008). Bean consumption is associated with greater nutrient intake, reduced systolic blood pressure, lower body weight, and a smaller waist circumference in adults: results from the National Health and Nutrition Examination Survey 1999-2002. *Journal of the American College of Nutrition*, 27(5), 569-576.
- Rabe, M., Ohri-Vachaspati, P., & Scheer, S. D. (2006). The influence of the youth expanded food and nutrition education program on nutrition knowledge and self-reported behaviors of elementary school children. *Journal of Extension* [On-line], 44(3). Available at: <http://www.joe.org/joe/2006june/rb6.php>
- Rhodes, D., Clemens, J., Goldman, J., Lacombe, R., & Moshfegh, A. (2012.) 2009-2010 *What we eat in America, NHANES Tables 1-36*. 2012. Retrieved from: <http://www.ars.usda.gov/Services/docs.htm?docid=18349>
- Roth-Yousey, L., Caskey, M., May, J., & Reicks, S. (2007). Modifying beverage choices of preadolescents through school-based nutrition education. *Journal of Extension* [On-line], 45(3). Available at: <http://www.joe.org/jpe/2007june/rb7.php>
- SAS Institute (2009). *SAS User's Guide. Statistics*, Version 9.2. SAS Inst. Inc., Cary, NC.
- Writing Group for the SEARCH for Diabetes in Youth Study Group, Dabelea, D., Bell, R. A., D'Agostino, R. B. Jr., Imperatore, G., Johansen, J. M., Linder, B., Liu, L. L., Loots, B., Marcovina, S., Mayer-Davis, E. J., Pettitt, D. J., & Waitzfelder, B. (2007). Incidence of diabetes in youth in the United States. *The Journal of the American Medical Association*, 297(24), 2716-2724. Retrieved from: <http://jama.jamanetwork.com/article.aspx?articleid=207653>

U. S. Department of Agriculture (USDA), & U.S. Department of Health and Human Services (HHS) (2010). *Dietary Guidelines for Americans, 2010*. 7th Edition; U.S. Government Printing Office; Washington, DC. Retrieved from:
<http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2010/PolicyDoc/PolicyDoc.pdf>

Copyright © by *Extension Journal, Inc.* ISSN 1077-5315. Articles appearing in the Journal become the property of the Journal. Single copies of articles may be reproduced in electronic or print form for use in educational or training activities. Inclusion of articles in other publications, electronic sources, or systematic large-scale distribution may be done only with prior electronic or written permission of the Journal Editorial Office, joe-ed@joe.org.

If you have difficulties viewing or printing this page, please contact [JOE Technical Support](#)