

12-2014

# Adaptation of Soil Judging to Northeast China

He Yun

*Clemson University*

Follow this and additional works at: [https://tigerprints.clemson.edu/all\\_theses](https://tigerprints.clemson.edu/all_theses)

 Part of the [Forest Sciences Commons](#)

---

## Recommended Citation

Yun, He, "Adaptation of Soil Judging to Northeast China" (2014). *All Theses*. 2076.

[https://tigerprints.clemson.edu/all\\_theses/2076](https://tigerprints.clemson.edu/all_theses/2076)

This Thesis is brought to you for free and open access by the Theses at TigerPrints. It has been accepted for inclusion in All Theses by an authorized administrator of TigerPrints. For more information, please contact [kokeefe@clemson.edu](mailto:kokeefe@clemson.edu).

ADAPTATION OF SOIL JUDGING TO NORTHEAST CHINA

---

A Thesis  
Presented to  
the Graduate School of  
Clemson University

---

In Partial Fulfillment  
of the Requirements for the Degree  
Master of Science  
Forest Resource.

---

by  
He Yun  
December 2014

---

Accepted by:  
Dr. Elena A. Mikhailova, Committee Chair  
Dr. Christopher J. Post  
Dr. Lawrence R. Gering

## ABSTRACT

Soil Judging teaches students important skills for field identification of soil types, properties, and interpretations for use. The adaptation of Soil Judging in Northeast China can be beneficial to students as well as government agencies and the private sector. The objective of this study was to adapt Soil Judging to the Northeast region of China by a graduate student from China, who was trained using an undergraduate course in Soil Judging and a regional Soil Judging competition. Unlike the U.S., China has 14 soil orders, with six soil orders somewhat similar to the ones found in the Southeast region of the U.S. A Southeastern Region Soil Judging Handbook was used for newly developed teaching materials for Northeast of China (including tables of soil physical and chemical properties, topographic maps, and scorecards). These new teaching materials can significantly improve soil education and mitigate problems associated with land use management.

## DEDICATION

To my mother Hongbo Fu, my dad Guoqing Yun and uncle Wenpu Liu for your love and support. Especially thank to my mother for everything over these years.

## ACKNOWLEDGMENTS

I wish to thank and acknowledge the students from FNR 4700: Soil Judging (Fall, 2013) course and organizers of the 2013 Southeast Regional Collegiate Soils Contest in Tennessee for helping to develop this project. Financial support for this project was provided by the Clemson University.

## TABLE OF CONTENTS

	Page
TITLE PAGE .....	i
ABSTRACT .....	ii
DEDICATION .....	iii
ACKNOWLEDGMENTS .....	iv
CHAPTER	
I.    INTRODUCTION .....	1
II.   MATERIALS AND METHODS.....	5
Study Site .....	5
Equipment needed for Soil Judging in Northeast China.....	6
Laboratory analysis .....	6
Topographic maps.....	6
Courses background.....	6
Learning management system.....	7
III.  RESULTS AND DISCUSSION.....	8
IV.  CONCLUSIONS.....	11
REFERENCES .....	12
APPENDIX.....	16

# **CHAPTER ONE**

## **INTRODUCTION**

Soil classification is an important component in the exchange and advancement of soil knowledge worldwide. Field description and laboratory analysis results are a foundation of soil classification. Various attempts have been made to correlate Chinese Soil Taxonomy with U.S. classification, but the need to further study the comparability lower classification levels still remains (Shi et al., 2004). Field and laboratory analysis provide essential data for soil classification, and there is a need to compare the use of the soil data at the field level in China and the US. Soil Judging is used in many countries of the world (e.g. USA and Germany) to train soil scientists to describe, classify, and interpret soil for different uses. The first International Soil Judging Contest took place in June 2014 at the 20th World Congress of Soil Science in Korea.

Adaptation of soil judging education to China can improve this soil science knowledge exchange, and can potentially alleviate land use problems in China by educating students and planners about important soil properties related to land use such as: soil infiltration rate, hydraulic conductivity, available water, soil wetness class, and soil interpretations related to suitability for dwellings with basements, septic tank absorption field, and local roads and streets.

Soil infiltration rate is the velocity at which water enters the soil, and it is one of the most important soil physical characteristics. It affects soil water availability for plant use and groundwater recharge, and is related to soil erosion and water runoff (Lipiec et al., 2006). It is affected by soil texture in the surface horizon (A) and it depends on soil texture (e.g. rapid for sand, loamy sand, or sandy loam texture with soil organic carbon

more than 1.2%). Northeast China is a predominantly agricultural area. Tillage practices could greatly affect soil macro porosity and infiltration characteristics, which consequently affect water runoff and soil erosion (Kemper et al., 2012). However, years of conventional tillage has resulted in increasing soil erosion and degradation problems as well as decrease in soil infiltration rates (Liu et al., 2010).

Soil saturated hydraulic conductivity is a quantitative expression of soil ability to transmit water under a given hydraulic gradient in the subsurface horizons (Julià et al. 2004). It is affected by soil texture, rock fragment, presence of restrictive layers (e.g. fragipan), structure, presence of redoximorphic depletions and color in the surface horizons (e.g. high, moderate, low). Soil texture is important for determining septic tank absorption field suitability (e.g. slight, moderate, or severe limitation). For example, Han et al. (2009), reported that a thick subsurface plow pan lowered the soil saturated hydraulic conductivity and additions of straw and organic manure resulted in increased saturated hydraulic conductivity. In another study by Zhang et al. (2013), high leachate levels of municipal solid waste were measured in the landfills in Southern China. Currently, ninety percent of villages in China don't have enough drainage channels and waste water treatments facilities, which results in the pollution of drinking water and the aquatic environment (Dong et al., 2012).

Available soil water capacity is the water held between field capacity and permanent wilting point and it is calculated to a depth of 150 cm of the soil or other thickness if the soil horizons are unfavorable for roots (Karathanasis et al., 2013). The available water calculations factors are (cm of water/cm of soil): 0.05 for sands, loamy sands; 0.15 for textures not included in other classes, and 0.20 for silt loam, silt, and silty

clay loam (Karathanasis et al., 2013). Zhou et al. (2005) note that laboratory-determined data for the available soil water capacity (ASWC) are often lacking in soil profiles databases in China. Geospatial analysis of ASWC showed the following pattern: higher ASWC values in the east compared to the Western part of China, and ASWC values are reduced from south to the north, except for the Northeastern part of China (Zhou et al., 2005).

Soil wetness class is the rate at which water is removed from the soil by both runoff and percolation and it is influenced by landscape position, slope gradient, infiltration rate, surface runoff, and permeability (Karathanasis et al., 2013). Redoximorphic features and soil color are used to determine the soil wetness class (Karathanasis et al., 2013). Ma and Fu (2006) reported evidence of drying trend over Northern China from 1951 to 2004. Topographic wetness index was highly correlated with soil organic matter content (Pei et al., 2010).

Suitability for dwellings with basements is affected by the following factors: flooding or ponding frequency, slope (%), depth to seasonally high water table, depth to soft rock “saprolite” (Cr) or depth to hard rock, cm (R) (Karathanasis et al., 2013). The degree of limitation varies from slight to moderate or severe (Karathanasis et al., 2013). Urbanization is widespread in China with the largest expansions in the Eastern Plain and the southeast (Liu and Tian, 2010).

Septic tank absorption field suitability is affected by the following factors: flooding or ponding frequency, slope (%), depth to seasonally high water table, limiting hydraulic conductivity, depth to soft rock “saprolite” (Cr) or depth to hard rock, cm (R) (Karathanasis et al., 2013). The degree of limitation varies from slight to moderate or

severe (Karathanasis et al., 2013). Septic tanks are used in many parts of the country, but there is a lack of appropriate waste water collection and treatment facilities in the rural areas (Chen et al., 2014). Lu et al. (2008) reported groundwater contamination from poor management of septic tanks in the southern coastal area of China, but the description of the study area raises many questions about septic tank absorption field suitabilities in the first place due to shallow water table, highly permeable soils, and frequent floods (especially in the areas with impervious surfaces).

Suitability for local roads and streets is affected by the following factors: flooding or ponding frequency, slope (%), depth to seasonally high water table, depth to soft rock “saprolite” (Cr) or depth to hard rock, cm (R) (Karathanasis et al., 2013). The degree of limitation varies from slight to moderate or severe (Karathanasis et al., 2013). Soil properties have significant impact on suitability for roads and streets in China as demonstrated by Du et al. (1999) by examining shrink-swell potential of natural and disturbed soils. Another study, by Fu et al. (2012), assessed land resources carrying capacity in Fujian Province of East China by using ecological suitability analysis in geographic information systems (GIS), and selecting altitude, slope, land use type, distance from resident land, distance from main traffic roads, and distance from environmentally sensitive area as the sensitive factors. Increased pressure on soil resources from urbanization and agriculture was also reported by Li (2012) in the context of urban-rural interactions patterns and dynamic land-use.

The objectives of this study were to develop a scorecard and recommendations for potential adaptation of Soil Judging education to the Northeast region of China.

## CHAPTER TWO

### MATERIALS AND METHODS

#### **Study Site**

Northeast China, is one of China's geographic and economic regions and it is composed of Heilongjiang, Jilin and Liaoning Provinces (Fig. 1, Table I). The largest plains in China are located in the Northeast region, which is rich in natural resources, culture, and economic development. It has a total surface area of about 1,370,000 km<sup>2</sup> and a population of about 121,000,000 (China Statistical Yearbook, 2012). The Northeast region has a semi-humid climate with an annual precipitation from 500 to 600 mm, with 90% of the precipitation falls as rain between April and September. It includes two of six ecological regions of China: the cool, subhumid Dongbei Plain (natural vegetation is a forest-steppe or meadow-steppe), and the cool, subhumid Da Hinggan (taiga forest), Xiao Hinggan and Changbai Mountain ranges (mixed needle- and broad-leaved forests) in the Northeast (ISSAS and ISRIC, 1994).

Soil Judging requires excavation of soil profiles, setting up slope stakes, scorecards (Fig. 2, 3), tables of soil physical and chemical properties (Fig. 4), and a set of soil judging equipment (Table IV). Score cards used for grading in soil judging competitions must be adapted to local soils and classification. Training for a 2013 soil judging competition at Tennessee Tech University enabled a graduate student from Northeast China to become familiar with several of the soil orders in Northeast China that are also present in the Southeastern United States (e.g. Alfisols, Entisols, and Inceptisols).

## **Equipment needed for Soil Judging in Northeast China**

A set of tools that must be provided for each student involved in a soil judging completion includes: a score card, official rules, an abney level or clinometer, garden spade, bucket, clip-board, soil collection trays, water bottle, measuring tape, a calculator, a pencil, and a Mussel color chart (Table IV). All items needed for a soil judging kit can be purchased locally in Northeast China (Table IV).

## **Laboratory analysis**

Soil samples to provide soil physical and chemical data for the students could be analyzed in any one of the several soil nutrient analysis laboratories in Northeast China: Northeast University Institute, and Liaoning Provincial Institute of Meteorology.

## **Topographic maps**

Soil profile locations were mapped as Geographic Information Systems (GIS) point shape-file files using ArcMap 10.2. Digital elevation models (DEM) surrounding each soil profile location (at a 30m resolution) were acquired from the ASTER digital elevation map (<http://asterweb.jpl.nasa.gov/gdem.asp>). Contour maps were created using the contour tool in ArcMap to create 30m topographic maps over the extent of the DEM.

## **Courses background**

Soil judging course can be incorporated in various soil science courses currently taught in Northeast China, for example: Jilin University (Environmental Soil Science); Heilongjiang University (Soil Science); Harbin Institute of Technology (Soil Science and

Soil Mechanics), and Liaoning Agriculture University (Soil Geography, Land Resource Science). Sectors that can also potentially benefit directly from soil judging education in China are the agriculture, housing and town planning, transportation, and health services.

### **Learning management system**

Modular Object-Oriented Dynamic Learning Environment (Moodle) is a free source e-learning software platform, which can be used for storing course materials, and assessing student' s learning via electronic quizzes and tests (<https://moodle.org/>).

Computer laboratories and ready access to the internet are available in most universities in the Northeast China.

## **CHAPTER THREE**

### **RESULTS AND DISCUSSION**

There three versions of Chinese soil classifications:

- Chinese Soil Taxonomic Classification System, 1st Proposal (1991), which distinguished 13 soil orders (Primarosols, Vertisols, Aridisols, Isohumols, Spodisols, Siallisols, Ferrallisols, Fersiallisols, Aquisols, Halosols, Anthrosols, and Andisols).

- Chinese Soil Taxonomic Classification System, Revised Proposal (1995), which distinguished 12 soil orders (Ferrasols, Alfisols, Semi-alfisols, Pedocals, Aridisols, Desert soils, Amorphic soils, Semi-aqueous soils, Aquous soils, Alkali-saline soils, Anthrosols, and Alpine soils).

- In 2001, China adapted “3rd” classification system with 14 soil orders: Histosols, Anthrosols, Spodosols, Andosols, Ferralosols, Vertosols, Aridosols, Halosols, Gleyosols, Isohumosols, Ferrosols, Argosols, Cambosols, Primosols (Zitong and Ganlin, 2007) with five soil orders (Argosols, Anthrosols, Isohumosols, Halosols and Histosols) found in the Northeast region of China (Table II, Table III) and correlated with soil taxonomy (ST) by Shi et al. (2004).

Newly developed soil judging scorecard (Fig. 2) is adapted for the third soil classification system (2001), but it can be adapted to previous classifications as well. Comparison of Chinese and U.S. soil field description and classification revealed common and dissimilar items used in the newly adapted scorecard (Table III). In Part A. Morphology of the scorecard the following items were different in Chinese classification compared to the US classification: prefix, master horizon abbreviations, boundary distinctness thicknesses, and structure. Part B. Soil Profile and Interpretations of the

scorecard is a new addition to the Chinese Soil Classification System since no comparable tables were found. In Part C. Site Characteristics of the scorecard the following items were different in Chinese classification compared to the US classification: position of the site, soil slope intervals, surface runoff and erosion potential (determined by experiment in China). In Part D. Soil Classification of the scorecard the following items were different in Chinese classification compared to the US classification: Epipedons, subsurface horizons, and soil orders.

In order to demonstrate how to use soil judging scorecard, soil profile CN 34 (ISSAS and ISRIC, 1994) was used to fill out the “practice” soil scorecard (Fig. 1, 2, 3) with markings in red color, which provides user with answers (Mikhailova and Post, 2014). In addition to soil judging scorecard, other supplemental materials are used: 1) Soil physical and chemical properties (Fig. 4), 2) optional topographic map of the area (Fig. 5), 3) textural triangle (not shown, but it is the same used in both China and USA), 4) abbreviations of distinctness of soil boundary, texture, modifiers of rock fragment quantity and size, structure grade, structure shape, consistence, redoximorphic features (Tables III, and IV), 5) tables of surface and soil erosion potential classes (Table VI), and 6) tables of soil use interpretations for dwellings with basement, septic absorption fields, and local roads and streets (Table V),. Soil profile CN 34 is one of the reference soil profiles of the People’ s Republic of China published in ISIS 4.0 data sheet. It has been classified as Silti-Chromic Cambisol (Eutric) in FAO/UNESCO (1988) classification, and as Typic Ustochrept, fine-silty, mixee, mesic in the USDA/SCS Soil Taxonomy (1992), and as Haplic cinnamon soil in Chinese Soil Taxonomy and Classification (1991), and as Cambisol in Chinese Soil Taxonomy and Classification (2001). The ISIS 4.0 data sheet

for CN 34 was used to fill out the scorecard. In Part B, infiltration rate was determined to be medium based on soil texture (SiL/SiCL) and soil organic carbon content (0.7%) in the Ap horizon (Karathanasis et al., 2013). Hydraulic conductivity was determined to be medium based on subsurface horizon characteristics (Karathanasis et al., 2013).

Available water was calculated based on depth of 150 cm x 0.20 (multiplier for SiL and SiCL in all of the horizons) (Karathanasis et al., 2013).. Soil wetness class is > 150 cm (not wet at depths of less than 151 cm) based on lack of redoximorphic features through the soil profile (Karathanasis et al., 2013). Soil interpretation for dwellings with basements, septic tank absorption fields, and local roads and streets was “2 = moderate” based on Table V using the following criteria: Flooding or ponding (None), Fig. 4, Slope (7-15%), depth to seasonally high water table and depth to soft rock “saprolite” , Cr (cm) > 100 cm, and depth to hard rock, R (cm) > 150 cm. In this case, slope (7-15%) was the moderate degree of limitation for all three uses. In Part C, surface runoff class was “rapid” based on 10% slope and “medium” infiltration in the Part B of the scorecard (Table VI). In Part C, erosion potential was “very high” based “rapid” surface runoff and SiL/SiCL surface horizon texture in the Part A of the scorecard (Table VI).

In general, U.S. scorecard can be used in China with necessary modifications depending on the region and soil interpretations to be used. The scorecard can further adapted to simultaneously train the user to describe and classify soil in multiple soil classifications.

## **CHAPTER FOUR**

### **CONCLUSIONS**

The introduction of soil judging to Northeast China may serve as a low cost, nontraditional way of teaching students and government workers simple and efficient techniques of land management and use. Northeast China has the basic infrastructure for a successful introduce of soil judging competitions to schools (middle and high schools, colleges, and universities), and various government sectors such as agriculture, health, road construction and building and town planning sectors. Soil nutrient analysis data can be obtained from any of the soil nutrient analysis laboratories in Northeast China.

## REFERENCES

- Chen, Z., Wen, Q., Guan, H., Bakke, R. and Ren, N. 2014. Anaerobic treatment of domestic sewage in modified septic tanks at low temperature. *Environ. Tech.* **35**: 2123-2131.
- Chen, X. 2002. The land resource and land use of Hong Kong. *Int. J. Remote. Sens.* **23**: 107-124.
- Dong, H., Qiang, Z., Wang, W. and Jin, H. 2012. Evaluation of rural wastewater treatments in a county of eastern China. *J. Env. Monit.* **14**: 1906-1913.
- Du, Y., Li, S. and Hayashi, S. 1999. Swelling-shrinkage properties and soil improvement of compacted expansive soil, Ning-Liang Highway, China. *Eng. Geol.* **53**: 351-358.
- Duan, X., Xie, Y., Liu, B., Liu, G., Feng, Y. and Gao, X. 2012. Soil loss tolerance in the black soil region of Northeast China. *J. Geogr. Sci.* **22**: 737-751.
- Fu, P. A. 2007. Variation of temperature, moisture and hydraulic conductivity of black soil in northeast of china. *Beijing Normal University (People's Republic of China)*.
- Fu, S., Zhang, P. and Jiang, J. 2012. Evaluation of land resources carrying capacity of development zone based on planning environment impact assessment. *J. Appl. Ecol.* (Chinese) **23**: 459-67.
- Gómez-Robledo, L., López-Ruiz, N., Melgosaa, M., Palmab, A. J., Capitán-Vallvey, L. F. and Sánchez-Marañón, M. 2013. Using the mobile phone as Munsell soil-colour sensor: An experiment under controlled illumination conditions. *Comput. Electron. Agr.* **99**: 200-208.
- Guo, Z. and Zhang, G. 2007. Chinese soil taxonomy: A milestone of soil classification in China. *Sci. Found. China.* **15**: 41-45.

- Han, X., Zou, W., Wang, F. and Wang, F. 2009. Construction effect of fertile cultivated layer in black soil. *Chinese J. Appl. Ecol.* **20**: 2996-3002.
- Institute of Soil Science. 1991. Chinese Soil Taxonomic Classification (1<sup>st</sup> proposal). Science Press, Beijing, China (In Chinese).
- Institute of Soil Science. 1995. Chinese Soil Taxonomy (revised scheme). China Agricultural Science and Technology Press, Beijing, China (In Chinese).
- Institute of Soil Science. 1995. Chinese Soil Taxonomy (3rd scheme). China Agricultural Science and Technology Press, Beijing, China (In Chinese).
- ISSAS and ISRIC. 1994. Reference soil profiles of the People's Republic of China. Field and analytical data. Country Report 2. (Gong Zitong, Luo Guobao, Zhang Ganlin, O.C. Spaargaren and J.H. Kauffman, compilers).
- Karathanasis, A.D., Galbraith, J.M., Shaw, J.N. and Thompson, J.A. 2013. Handbook for Collegiate Soils Contest. Southeast Region Web Site:  
[http://gis.clemson.edu/elena/documents/SE\\_Handbook\\_2013.pdf](http://gis.clemson.edu/elena/documents/SE_Handbook_2013.pdf)
- Lee, L. 1999. Labor location, conservation, and land quality: The case of West Jilin, China. *Ann. Assoc. Am. Geogr.* **89**: 633–657.
- Li, X., Liu, L., Wang, Y., Luo, G., Chen, X., Yang, X., Hall, H. P. M., Guo, R., Wang, H., Cui, J. and He, X. 2013. Heavy metal contamination of urban soil in an old industrial city (Shenyang) in Northeast China. *Geoderma* **192**: 50–58.
- Li, Y. 2012. Urban-rural interaction patterns and dynamic land use: implications for urban-rural integration in China. *Reg. Environ. Change.* **12**: 803–812.
- Liu, X., Zhang, S., Zhang, X., Ding, G. and Cruse, R. M. 2011. Soil erosion control practices in Northeast China: A mini-review. *Soil Till. Res.* **117**: 44.

- Liu, X.B., Zhang, X.Y., Wang, Y.X., Sui, Y.Y., Zhang, S.L., Herbert, S. J. and Ding, G.W. 2010. Soil degradation: a problem threatening the sustainable development of agriculture in Northeast China. *Plant Soil Environ.* **56(2)**: 87–97.
- Liu, M. and Tian, H. 2010. China's land cover and land use change from 1700 to 2005: Estimations from high-resolution satellite data and historical archives. *Global Biogeo. Cycles.* 24: GB3003.
- Lu, Y., Tang, C., Chen, J. and Sakura, Y. 2008. Impact of septic tank systems on local groundwater quality and water supply in the Pearl River Delta, China: case study. *Hydrol. Process.* **22**, 443-450.
- Ma, Z. and Fu, C. 2006. Some evidence of drying trend over northern China from 1951 to 2004. *Chinese Sci. Bulletin.* **51**: 2913-2925.
- Mikhailova, E.A. and Post, C.J. 2014. Laboratory Exercises for Soil Judging. 2<sup>nd</sup> edition. Printed in the United States of America.
- Pei, T., Qin, C.Z., Zhu, A.X., Yang, L., Luo, M., Li, B.L. and Zhou, C.H. 2010. Mapping soil organic matter using the topographic wetness index: A comparative study based on different flow-direction algorithms and kriging methods. *Ecol. Ind.* **10**: 610-619.
- Qi, S. and Luo, F. 2006. Land-use change and its environmental impact in the Heihe River Basin, arid northwestern China. *Env.. Geol.* **50**: 535–540.
- Shi, X., Yu, D., Sun, W., Wang, H., Zhao, Q. and Gong, Z. 2004. Reference benchmarks relating to great groups of genetic soil classification of China with soil taxonomy. *Chinese Sci. Bulletin.* **49**: 1507-1511.

- Soil Survey Staff. 1999. Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys. 2<sup>nd</sup> edition. USDA/NRCS. Agriculture Handbook No. 436.
- Soil Survey Staff. 2003. Natural Resources Conservation Service, United States Department of Agriculture. Keys to Soil Taxonomy. 9th ed. USDA. Washington, DC.
- Xu, S. G., Xu, X. Z., Xu, Y., Chen, S. C. and Zhang, H. W. 2010. Soil loss and conservation in the black soil region of Northeast China: a retrospective study. *Environ. Sci. Pol.* **13**: 793–800.
- Zhang, W. J., Zhang, G. G. and Che, Y. M. 2013. Analyses on a high leachate mound in a landfill of municipal solid waste in China. *Env. Earth Sci.* **70**: 1747-1752.
- Zhang, X. Y., Cruse, R. M., Sui, Y. Y. and Jhao, Z. 2006. Soil compaction induced by small tractor traffic in Northeast China. *Soil Sci. Soc. Am. J.* **70**: 613.
- Zhao, M. and Zeng, X. 2002. A theoretical analysis on the local climate change induced by the change of landuse. *Adv. Atmos. Sci.* **19**: 45–63.
- Zhou, W., Liu, G., Pan, J. and Feng, X. 2005. Distribution of available soil water capacity in China. *J. Geogr. Sci.* **15**: 3-12.
- Zhu, H., Lu, C. and Li, X. 2004. Land use studies in China. *J. Geogr. Sci.* **14**: 69–73.

## APPENDIX





Fig. 1. Map of Northeast of China with soil pit locations (adapted from ISSAS and ISRIC, 1994).

### 中国地区大学生土壤大赛

地点 34  
 土层数 3  
 研究土壤深度 150 cm  
 第三土层钉子深度 80 cm

- 质地  
 1. 不成球=沙子 (0-10%粘土含量)  
 2. 球破碎=壤沙土(0-15%粘土含量)  
 3. 土壤条<1英寸(粘性) 0-27%粘土含量  
 4. 土壤条1-2英寸(粘壤土), 20-40%粘土含量  
 5. 土壤条>2英寸(粘土), 40-100% 粘土含量

姓名 \_\_\_\_\_

一、形态

土层				质地				颜色			结构		组成			土壤氧化物			分数
前缀	符号	附加符号	编号	最低深度	过渡程	岩屑丰	质地划分	粘土含量	色相	明度	纯度	发育程	形状	结持性	强氧化	氧化	衰减基		
1	3	2	1	3	1	1	3	1	1	1	1	2	1	1	1	1	1	1	27
—	A	p	—	15	A	—	SiL/SiCL	29	7.5YR	4	6	B	H	B	无	无	无		
—	B	w	1	65	B	—	SiCL	33	5YR	3	4	C	F	C	无	无	无		
—	B	w	2	150+	—	—	SiCL	33	5YR	4	8	C	I	B	无	无	无		

注意: 如果表格里存在R层或者Cr层不用具体考虑他们的性质, 只在表格中填入破折号(-)。如果R层在Cr层下面也不用考虑Cr层

二、土壤剖面及说明

<p><b>渗透速率 (5)</b> (在一个土层中)</p> <p>急 (S, LS; or SL with &gt;2%OM)</p> <p>X 中 (所有其他)</p> <p>慢 (C, SiC, SC &amp; massive &amp; weak str. Or dense layers, R)</p> <p><b>土壤饱和导水率 (5)</b></p> <p>高 (第二层土层: S, LS, w/out GR)</p> <p>X 中 (其他)</p> <p>低 (多个或者最少有一个土层: C, SC, SiC并且大块、弱结构或者板块的, 或者氧化匮乏。颜色等级&gt;4 色度≤2)</p>	<p>(0.05 = S, LS; 0.15 =其他, 0.20 = SiL, Si, SiCL)</p> <p><b>潜水深度 (5)</b></p> <p>非常低 ≤ 7.5 cm</p> <p>低 &gt; 7.5 and ≤ 15.0 cm</p> <p>中 &gt; 15.0 and ≤ 22.5 cm</p> <p>X 高 &gt; 22.5 cm (150x0.2=30 cm)</p> <p><b>土壤湿润等级 (5)</b></p> <p>X &gt; 150 cm (1) 最顶上的A层含有:</p> <p>101-150 cm a) 土壤色调≤2且</p> <p>51-100 cm b) 不同的氧化匮乏或者衰减基质</p> <p>25-50 cm c) 氧化匮乏或者衰减基质直接在A土层下</p> <p>&lt; 25 cm (2) 可以观察到的颜色等级≥4且色度≤2的最浅深度</p>	<p>(根据提供的表格可轻松判定)</p> <p><b>土地应用</b></p> <p>2 带地下室的住房</p> <p>2 化粪池吸收场所</p> <p>2 街道</p> <p>(1 = 轻微, 2 = 中度, 3 = 重度 局限性)</p>
第一部分 _____		
第二部分 _____		
第三部分 _____		
第四部分 _____		
总分 _____		

Fig. 2. Front side of scorecard for Northeast China Soil Contest (adapted from Karathanasis et al., 2011). Text in red color is a teaching-aid for practice in the field. During Soil Judging competitions, the red text is not used.

三、采样点特征

采样点地形 (5)

- 凹地
- 河道
- 河滩
- 山麓
- 河成阶地
- 高地

成土母质 (5)

- 冲积物(河流)
- 崩积物(受重力影响多见于山地)
- 其他母质(由当地形成)

坡度(5)

- 平坡(0--3%)
- 微坡(3--7%)
- 缓坡(7--15%)
- 中坡(15--25%)
- 陡坡(25--35%)
- 极陡坡(>35%)

地表径流 (5)

(表格见后, 请学会使用)

- 池塘
- 非常慢
- 慢
- 中等
- 快
- 非常快

侵蚀势 (5)

(表格见后, 请学会使用)

- 非常低
- 低
- 中等
- 高
- 非常高

四、土壤分类

表层 (5)

- 暗脊表层
- 淡薄表层
- 暗沃表层
- 肥熟表层
- 盐积层

第二土层和其性质 (5分每点)

- 白土层
- 锥形层
- 黏磐
- 耕作淀基层
- 粘化层
- 碱基层
- 盐积层
- 无

土纲(5) (中国土壤系统分类第三版2001)

- 淋溶土
- 均腐土
- 人为土
- 盐成土
- 有机土
- 锥形土

第三部分得分 \_\_\_\_\_

第四部分得分 \_\_\_\_\_

Fig. 3. Back side of scorecard for Northeast China Soil Contest (adapted from Karathanasis et al., 2011). Text in red color is a teaching-aid for practice in the field. During Soil Judging competitions, the red text is not used.

# PIT 34

No. of horizons 3

Depth to be described 150 cm

Nail in the 3<sup>rd</sup> horizon @ 80 cm

HORIZON	OC (%)	BS (%)
1	0.7	23
2	0.3	52
3	0.2	44

Flooding: NONE

Ponding: NONE

Fig. 4. Soil physical and chemical properties for the soil profile No. 34 in Liaoning Province, Chaoyang, China (adapted from ISSAS and ISRIC, 1994).





Fig. 5. Topographic map for the soil profile No. 34 in Liaoning Province, Chaoyang, China (adapted from ISSAS and ISRIC, 1994).




Fig. 6. Examining a no-pick zone of the practice soil profile during 2013 Southeastern Regional Soil Judging Competition in TN Tech University, TN.



Fig. 7. Slope determination in the field.



Fig. 8. Texture by feel during 2013 Southeastern Regional Soil Judging Competition in TN Tech University, TN.



Fig. 9. Clemson University Soil Judging team (He Yun, graduate student from China is far right) 2013 Southeastern Regional Soil Judging Competition in TN Tech University, TN.

Table I. General information of the Northeast of China(Source: China Statistical Yearbook, 2012).

Province	Capital	Population (2010)	Area km <sup>2</sup>	Population density people/km <sup>2</sup>	Land-use
Heilongjiang	Harbin	38,312,224	454,000	84.38	Logging, timber, industry, farming
Jilin	Changchun	27,462,297	187,400	146.54	Logging, timber, farming
Liaoning	Shenyang	43,745,323	145,900	299.83	Farming. industry,

TABLE II. Soils of the Northeast China according to 2001 Chinese Classification System.

Heilongjiang	Jilin	Liaoning
Chinese Classification, 3 <sup>rd</sup> proposal (2001)		
Argosols (淋溶土)	Argosols (淋溶土)	Argosols (淋溶土)
Isohumosols (均腐土)	Isohumosols (均腐土)	Isohumosols (均腐土)
Halosols (盐成土)	Halosols (盐成土)	Anthrosols (人为土)
Histosols (有机土)		Halosols (盐成土)
Cambisols (雏形土)		Cambisols (雏形土)

TABLE III. Comparison of Chinese and U.S. soil description and classification used in the soil judging scorecard based on Southeast region of USA and Northeast region of China.

Scorecard	U.S. (Source: Soil Survey Staff, 1999)	China (Source: Institute of Soil Science, 1991)
<b>A. Morphology</b>		
Prefix	1, 2...	-
Master horizon abbreviation	O, A, E, B, C, R	O, K, A, E, B, G, C, R
Horizon Sub	a, b, c, g, h, k, m, n, p, q...	a, b, c, g, h, k, m, n, p, q...
No.	1, 2, 3	-
Lower depth	cm	cm
Boundary distinction	A (<2cm), C (2-6cm), G (6-16cm), D (>16cm)	A(<2cm), B(2-5cm), C(5-12cm), D(>12cm)
Rock fragment	Percent by volume	Percent by volume and the rock size
USDA class	Texture-by-feel (Textural triangle)	Texture-by-feel (Textural triangle)
Clay content	Particle size, texture-by-feel	Particle size, texture-by-feel
Color	Munsell Color Chart	Munsell Color Chart Describe by words
Structure grade	SLS, WK, MO, ST	SLS, WK, MO, ST
Structure shape	GR, PL, MA, PR, SG, ABK, SBK	Platy, Scaly, Prismatic, Columnar, Prismatic Blocky, Lumpy, Core shape, Granular, Pellets, Crumb
Consistency (moist)	L, Fi, FR, VFi, VFR, EFi	L, Fi, FR, VFi, VFR, EFi
<b>B. Soil profile and Interpretations</b>		

Infiltration rate	Determined from scorecard	By experiment
Hydraulic conductivity	Determined from scorecard	By experiment
Available water	Determined from scorecard	By experiment
Soil wetness class	Determined from scorecard	-
Soil interpretations	Determined from scorecard tables	-

### C. Site characteristics

Position of site	Depression, Drainage Way, Flood Plain, Footslope, Stream Terrace, Upland (Field observations, topographic maps)	Mountain, Hills, Plain, Plateau, other
Parent material	Alluvium, Colluvium, Residuum	Alluvium, Colluvium, Residuum
Soil slope	Nearly Level(0-2), Gently Sloping(2-6), Sloping(6-12), Moderately Sloping(12-20), Strongly Sloping(20-30), Steep(>30)	<3, 3-7, 7-15, 15-25, 25-35, >35
Surface runoff	Ponded, Very Slow, Slow, Medium, Rapid, Very Rapid	By experiment
Erosion potential	Very Low, Low, Medium, High, Very High	By experiment

### D. Soil classification

Epipedons	Mollic, Ochric, Umbric	Mollic, Ochric, Umbric, Fimic, Salic
Subsurface horizons and characteristics	Albic, Argillic, Cambic, Fragipan, Lith. Discontinuity, Lithic Contact, Paralithic Contact, None	Albic, Cambic, Agric, Argic, Claypan, Alkalic, Salipan, None
Order	Based on data	Based on data

---

1 TABLE IV. Abbreviation list for Soil Judging.

2 缩写

3 来源: Adapted from Handbook for Collegiate Soils Contest, 2013.

土层过渡程度

突然过渡=A

明显过渡=B

逐渐过渡=C

模糊过渡=D

质地划分

砂土= S

粉土= Si

粘土= C

壤土=L

砂壤土= SL

粉壤土= SiL

粘壤土= CL

壤砂土=LS

砂粘壤土= SCL

粉黏壤土= SiCL

砂粘土= SC

粉粘土= SiC

岩屑大小和丰度

很小=A

小=B

中=C

大=D

很大=E

无=a

少=b

中=c

多=d

### 土壤发育程度

无结构=A	弱发育程度=B	中度发育程 度=C	强发育程度 =D
-------	---------	--------------	-------------

### 土壤结构形状

片状=A	鳞片状=B	棱柱状=C	柱状=D
棱块状=E	团块状=F	核状=G	粒状=H
团粒状=I	屑粒状=J		

### 结持性

松散=A	极疏松=B	疏松=C
坚实=D	很坚实=E	极坚实=F

### 土壤氧化物

发现的话填“有”，没有发现填“无”

4 TABLE V. Translation of soil use interpretations tables.

5 土壤应用诠释表格

6 来源: Adapted from Handbook for Collegiate Soils Contest, 2013.

7

8 带地下室的住宅: 编辑自 NSSH Table 620-3.

影响利用的因素	限制程度		
	轻微 (1)	中等 (2)	严重 (3)
1. 发生洪水或者积水的频率	从不	没有提供	很少或者频繁
2. 坡度(%)	< 6	6 - 20	> 20
3. 地下水位深度(cm)	> 100	50 - 100	< 50
4. 土层Cr深度	> 100	50 - 100	< 50
5. 土层R深度	> 150	100 - 150	< 100

9

10 化粪池吸收场所: 编辑自 NSSH Table 620-17.

影响利用的因素	限制程度		
	轻微 (1)	中等 (2)	严重 (3)
1. 发生洪水或者积水的频率	从不	没有提供	很少或者频繁
2. 坡度(%)	< 6	6 - 20	> 20
3. 地下水位深度(cm)	> 150	100 - 150	< 100
4. 水利传导系数	中等	没有提供	低或者高
5. 土层Cr或者R的深度	> 150	100 - 150	< 100

11

12 街道: 编辑自 NSSH Table 620-5.

影响利用的因素	限制程度		
	轻微 (1)	中等 (2)	严重 (3)
1. 发生洪水或者积水的频率	从不	没有提供	很少或者频繁
2. 坡度(%)	< 6	6 - 20	> 20
3. 地下水位深(cm)	> 50	25 - 50	< 25
4. 土层Cr深度	> 100	50 - 100	< 50
5. 土层R深度	> 150	100 - 150	< 100

13

14 TABLE VI. Translation of surface runoff classes and soil erosion potential classes tables.

15 地表径流等级

16 来源: Adapted from Handbook for Collegiate Soils Contest, 2013.

	渗透速率快	渗透速率中等	渗透速率缓慢
凹地	积水	积水	积水
坡度0-1%	非常慢	非常慢	慢
坡度>1-2%	非常慢	慢	中等
坡度>2-6%	慢	中等	快
坡度>6-12%	中等	快	非常快
坡度>12%	快	非常快	非常快

17

18 侵蚀势等级

19 Source: Handbook for Collegiate Soils Contest, 2013.

地表径流等级	第二层土层质地			
	S, LS	SCL, SC	SL, CL, C, SiC	L, Si, SiL, SiCL
积水	非常慢	非常慢	非常慢	非常慢
非常慢	非常慢	非常慢	慢	中等
慢	非常慢	慢	中等	中等
中等	非常慢	慢	中等	高
快	慢	中等	高	非常高
非常快	中等	高	非常高	非常高

Table VII. Soil Judging equipment cost per person in China.

No.	Equipment/supplies	Cost in US	Cost in China (RMB as of 1/30/14)
1.	Abney level, clinometers, or other hand level	1	150
2.	Knife	3.5	20
3.	Water bottle	0.5	3
4.	Munsell Color Chart	150	1000
5.	Scorecards and supplemental materials	0.5	1
6.	Calculator	2	10
7.	Mechanical pencil	0.5	2
8.	Measuring tape and nail	1	3
9.	Clip board	1	10
10.	Containers for soil samples	0.5	20
11.	Bucket	1	6
	Total	185.5	1225