

Clemson University

TigerPrints

All Theses

Theses

12-2022

Characterization of *Colletotrichum nymphaeae* Isolates From Apple with Reduced Sensitivity to Fluazinam and Tebuconazole

Rulyu Meng
rulyum@g.clemson.edu

Follow this and additional works at: https://tigerprints.clemson.edu/all_theses



Part of the [Plant Pathology Commons](#)

Recommended Citation

Meng, Rulyu, "Characterization of *Colletotrichum nymphaeae* Isolates From Apple with Reduced Sensitivity to Fluazinam and Tebuconazole" (2022). *All Theses*. 3908.

https://tigerprints.clemson.edu/all_theses/3908

This Thesis is brought to you for free and open access by the Theses at TigerPrints. It has been accepted for inclusion in All Theses by an authorized administrator of TigerPrints. For more information, please contact kokeefe@clemson.edu.

CHARACTERIZATION OF COLLETOTRICHUM NYMPHAEAE ISOLATES
FROM APPLE WITH REDUCED SENSITIVITY TO FLUAZINAM AND
TEBUCONAZOLE

A Thesis
Presented to
the Graduate School of
Clemson University

In Partial Fulfillment
of the Requirements for the Degree
Master of Science,
Plant Science

by
Rulyu Meng
December 2022

Accepted by:
Dr. Guido Schnabel, Committee Chair
Dr. Julia Kerrigan
Dr. James Faust

ABSTRACT

Apple bitter rot is caused by *Colletotrichum nymphaeae* and other *Colletotrichum* species and management relies primarily on synthetic pesticides. Very few fungicides are effective against the disease and resistance has further limited their usefulness. A recent study indicated relatively low EC₅₀ values (the concentration required to inhibit 50% of mycelial growth in vitro) of *C. nymphaeae* isolates from Brazilian apples to fluazinam and tebuconazole, two fungicides that are not routinely used for bitter rot control. Isolates on opposite sides of the sensitivity range were designated either sensitive (lowest EC₅₀ values) and reduced-sensitive (highest EC₅₀ values). The objective of this research was to confirm stability of EC₅₀ values after storage, assess fluazinam and tebuconazole for apple bitter rot management in detached fruit studies, and to sequence the fungicide's putative target genes. Our research confirmed stability of EC₅₀ values after 8 months of cold storage. Detached fruit studies revealed that protective treatment of cv. Gala apple fruit with fluazinam and tebuconazole controlled both sensitive and reduced-sensitive phenotypes but if used curatively (after infection) reduced-sensitive isolates were controlled less effectively. The complete *OS-1*, *CYP51A*, and *CYP51B* genes from *C. nymphaeae* isolates sensitive or reduced-sensitive to fluazinam (potential target *OS-1*) and tebuconazole (*CYP51A* and *CYP51B*) were sequenced and three nucleotide changes located in two codons of *OS-1* leading to three amino acid changes were identified in reduced-sensitive isolates. This study underscores the potential usefulness for fluazinam and tebuconazole for bitter

rot control of apple and provides the genetic basis for studies on potential resistance mechanisms.

ACKNOWLEDGEMENTS

I would like to thank Dr. Guido Schnabel to offer me the opportunity work with him during graduate school. I would also like to thank my committee members, Dr. Julia Kerrigan, and Dr. James Faust for their guidance during my time as a master's student. I also would like to thank for Mr. Zhigang Li for his guidance. I also would like to thank Dr. Jhulia Gelain and Melissa Muñoz for their help with statistical analysis. I especially would like to thank Mitchell D. Rich and William Gura for helping our molecular and fruit experiments.

TABLE OF CONTENTS

	Page
TITLE PAGE	i
ABSTRACT.....	ii
ACKNOWLEDGMENTS	iv
TABLE OF CONTENTS.....	v
LIST OF TABLES.....	vii
LIST OF FIGURES	vii
CHAPTER	
I. LITERATURE REVIEW	1
1.1 The <i>Colletotrichum</i> spp.....	1
1.2 Importance of apples to the Brazilian economy and main disease issues.....	2
1.3 Bitter rot of apple; optimal condition and species involved	2
1.4 Apple bitter rot disease cycle.....	3
1.5 Chemical control of apple bitter rot.....	4
II. CHARACTERIZATION OF <i>COLLETOTRICHUM NYMPHAEAE</i> ISOLATES FROM APPLE WITH REDUCED SENSITIVITY TO FLUAZINAM AND TEBUCONAZOLE.....	7
2.1 INTRODUCTION	7
2.2 MATERIALS AND METHODS.....	11
2.2.1 <i>Colletotrichum nymphaeae</i> isolates	11
2.2.2 In vitro sensitivity to fluazinam and tebuconazole	12
2.2.3 Fungicide sensitivity assays using detached fruit	12
2.2.4 DNA extraction.....	14
2.2.5 Primer design and PCR amplification.....	14
2.2.6 Sequence analysis	17
2.2.7 Statistical analysis.....	17

Table of Contents (Continued)

	Page
2.3 RESULTS	18
2.3.1 <i>Colletotrichum nymphaeae</i> sensitivity to fluazinam and tebuconazole.....	18
2.3.2 Detached apple fruit assay	19
2.3.3 Sequencing of the <i>OS-1</i> , <i>CYP51A</i> , and <i>CYP51B</i> genes	21
2.4 DISCUSSION	22
LITERATURE CITED	27

LIST OF TABLES

Table	Page
2.2.5 Primers used for amplification of <i>CYP51A</i> , <i>CYP51B</i> and <i>OS-1</i> gene	15
2.3.1 Isolate names, fungicide sensitivity phenotype, geographical origin, year of isolation and EC ₅₀ values obtained before and after long-term storage of <i>Colletotrichum nymphaeae</i> isolates from Brazilian apple orchards	19

LIST OF FIGURES

Figure	Page
2.2.5 <i>CYP51A</i> and <i>CYP51B</i> gene length, intron locations, and locations of primers used in this study.....	17
2.4.2 Protective and curative activity of (A) fluazinam (Flu) and (B) tebuconazole (Teb) against <i>Colletotrichum nymphaeae</i> isolates with different fungicide sensitivity phenotypes on detached apple fruit.....	21
2.4.3 <i>OS-1</i> gene length, intron locations, variable nucleotide sequence locations leading to amino acid changes, and locations of primers used in this study	22

CHAPTER ONE

1. LITERATURE REVIEW

1.1 The *Colletotrichum* spp.

Colletotrichum species cause diseases in a wide range of hosts (Freeman, 2008; Afanador-Kafuri et al., 2003; Eastburn and Gubler, 1990; Freeman et al., 2002). They affect a wide range of economically important crops, including apple, peach, grape, blueberry, cranberry, and strawberry (Bernstein et al., 1995; Daykin and Milholland, 1984; Howard, 1972). There are different names for diseases caused by *Colletotrichum* spp. depending on the host and plant part. Anthracnose is the most commonly used disease name, while apple bitter rot or grape ripe rot are also used to describe diseases caused by *Colletotrichum* spp on apple and grapes, respectively. Despite *Colletotrichum* being the eighth most important genus of phytopathogenic fungi in the world (Crouch et al., 2014; Dean et al., 2012), the systematics of *Colletotrichum* species has been difficult to establish (Hyde et al., 2009) because of changing taxonomy (Bernstein et al., 1995). Phylogenetic analysis of the genus *Colletotrichum* reveals that it is comprised of nine major clades, as well as several smaller clusters and isolated species (Cannon et al., 2012). Among the nine major clades of *Colletotrichum*, the *C. acutatum* species complex and the *C. gloeosporioides* species complex are the two most common clades containing species that cause bitter rot disease of apples (Jayawardena et al., 2016; Peres et al., 2005; Munir et al., 2016).

1.2 Importance of apples to the Brazilian economy and main disease issues

Apple (*Malus domestica* Borkh.), native to central Asia and then introduced to the west and other parts of the world (Dzhangaliev, 2002; Forsline et al., 2003), is a major fruit crop cultivated in temperate regions today (Khodadadi et al., 2020). In Brazil, four regions account for the bulk of apple production: Santa Catarina state, Rio Grande do Sul state, Parana state, and Sao Paulo state (Ming, 2019). The apples produced in Brazil amounted to 983,247 tons in the 2020 season (FAO, 2022). Apple crops are vulnerable to a wide range of diseases affecting yield and fruit quality (Khodadadi et al., 2020). The three most important fungal diseases occurring frequently in Brazilian apple orchards are apple bitter rot and Glomerella leaf spot, both caused by *Colletotrichum* species (Velho et al. 2015), and apple scab caused by *Venturia inaequalis* (Bogo et al., 2012).

1.3 Bitter rot of apple; optimal conditions and species involved

Apple bitter rot was first described in England in 1856 and later in the United States (Taylor, 1971). Disease development depends on wet and warm weather conditions (Carraro et al., 2022). *C. nymphaeae*, *C. paranaense*, and *C. melonis* species causing apple bitter rot symptoms on detached fruit had optimal conidia germination at temperatures 20-25°C (Moreira et al., 2020). Optimal mycelium growth temperatures of individual *Colletotrichum* spp. range from 22 to 30°C (Dowling et al., 2020; Wang et al., 2015). The ideal weather for apple bitter rot infections is furnished by high moisture (80-100%) with frequent rains (Roberts, 1918; Ivey, 2016). All currently

available commercial apple cultivars are susceptible to the pathogen (Denardi et al., 2003). In Brazil, the disease was first described in the 1970s when commercial apple production was initiated in the country (Denardi et al., 2003). *C. nymphaeae* and *C. fiorinae* from the *Colletotrichum acutatum* species complex and *C. fructicola*, *C. aenigma*, and *C. siamense* from the *Colletotrichum gloeosporioides* species complex are known to cause apple bitter rot worldwide (Weir et al., 2012; Jayawardena et al., 2016).

1.4 Apple bitter rot disease cycle

Apple bitter rot is defined by fruit infections, but the causal agent may infect other plant tissues including leaves, petals, twigs, and fruitlets (Hamada et al., 2020; Everett et al., 2018; Nekoduka et al., 2018; Peres et al., 2005). Apple bitter rot produces symptoms initially consisting of small brown spots with red halos (Abid et al., 2019). Thereafter, the lesions enlarge rapidly and become circular and sunken (Abid et al., 2019) and Conical regions of necrotic brown tissue develop beneath the lesions (Munir et al., 2016). Copious amounts of conidia are produced in fruiting bodies called acervuli. Under moist, humid conditions, the spore masses appear creamy and are orange to pink-colored and arranged concentrically (Beever et al., 1995; Oo et al., 2018).

Colletotrichum spp may overwinter in mummified fruit in the canopy or on the ground where they serve as important inoculum sources for the following season (Roberts, 1918). In spring, *C. acutatum* starts to sporulate in orange horn-like structures on mummies, cankers, or apple bud scales (Roberts, 1918; Everett et al., 2018; Børve

and Stensvand, 2007), and emerging spores may contribute to flower infections. Tree flowers from apple, cherry, and peach can harbor the pathogen in the form of quiescent infections until optimal conditions favor disease development (Hamada et al., 2020; Stensvand et al., 2017; Børve and Stensvand, 2007; Moral et al., 2009). *C. acutatum* produces spores in both spring and summer; conidia from acervuli are rain-splash dispersed, germinate to form appressoria (Everett et al., 2018), and quiescent infections as appressoria or hyphae are established within fruit surfaces (Sinclair, 1991; Timmer et al., 1998). In summer, the primary inoculum sources are infected decaying petals, bud scales, twigs, fruitlets, and twig cankers (Everett et al., 2018). Infections may remain latent in apple fruit until environmental conditions and fruit maturity favor symptom development (Jones et al., 1996).

1.5 Chemical control of apple bitter rot

Fungicides with multisite modes of action (MOA) such as captan and mancozeb as well as single-site MOA are used to control fungal diseases in apple production. Compared to single-site MOA fungicides, the multisites are typically less effective but broader in their activity spectrum (i.e. they work against pathogens of different genera). The most commonly used chemical classes with single site MOA for bitter rot control are the quinone outside inhibitors (QoIs) and benzimidazoles (MBC). *Colletotrichum* spp. have been reported to develop resistance to many fungicides, including QoIs and MBCs (Avila-Adame et al., 2003; Hu et al., 2015; Poti et al., 2020). QoI fungicides inhibit the cytochrome bc1 respiratory complex in the fungal

mitochondria (Kim et al., 2003). QoI resistance has already been reported in apple orchards, specifically in *Venturia inaequalis*, the causal agent of apple scab (Lesniak et al., 2011). Control failure due to QoI resistance in *C. acutatum* was first documented in Florida strawberries in the 2013-2014 season (Forcelini and Peres, 2018) and subsequently in *C. nymphaeae* isolates from Ohio strawberries (Chechi, et al. 2020). Resistance to QoI fungicides has also been described for isolates belonging to the *C. gloeosporioides* species complex and obtained from diseased apples (Martin et al., 2022; Chechi et al., 2020; Yokosawa et al., 2017). Due to the increasing occurrence of QoI resistance in *Colletotrichum* species as well as reports of resistance to MBC fungicides in the *C. gloeosporioides* complex (Hu et al., 2015), captan has become an important fungicide in resistance management and the management of common apple diseases such as apple scab (*Venturia inaequalis*), apple bitter rot (Abbott and Beckerman, 2018), and anthracnose of strawberry (MacKenzie and Peres, 2012).

Previous studies suggest that *C. nymphaeae* isolates, belonging to *C. acutatum* species complex, were highly resistant to MBCs but sensitive to DMIs (Gelain et al., in press). To the best of our knowledge, no field resistance to tebuconazole or fluazinam has been described. Because of the limited number of effective fungicides for bitter rot control in Brazil, chemicals with other MOA, such as fluazinam and tebuconazole, become more promising for the potential suppression of bitter rot.

Fluazinam is a single site MOA fungicide used in agriculture against various diseases. The mode of action involves the uncoupling of oxidative phosphorylation in

mitochondria (Guo et al., 1991) and high reactivity with thiols. Fluazinam is an aminopyridine inhibiting glucose oxidation in a dose-dependent manner and suppressing the energy production process through direct inhibition of the adenosine triphosphate (ATP) synthetase in fungal cells (Vitoratos, 2014). Based on recent studies, fluazinam successfully controlled strawberry crown rot caused by *C. gloeosporioides* (Oliveira et al., 2020) and seed decay by *C. truncatum* (Oh and Kang, 2002). Tebuconazole is a demethylation inhibitor (DMI) fungicide that targets the sterol 14 α -demethylase (Chen et al., 2018) and has become an integral part of apple scab management (Ilhan et al., 2006; Arslan, 2016). Resistance mechanisms include modifications or overexpression of target genes. For example, DMI fungicide resistance in *Colletotrichum truncatum* was linked to *CYP51A* and *CYP51B* target gene expression (Chen et al., 2018). According to Oliveira et al. (2020) and Silva-Junior et al. (2014), tebuconazole can inhibit strawberry crown rot caused by *C. gloeosporioides* and citrus anthracnose caused by *C. acutatum*.

Conventional fungicide applications in apple production are critical for disease management. However, the registration of new fungicides is hindered due to cost of product development and due to difficulty in new product registration. The use of existing products with suppressive action may therefore be beneficial at least in the short term for IPM (Integrated Pest Management) practices.

CHAPTER TWO

2.1 INTRODUCTION

Apples (*Malus domestica* Borkh.) are an economically important fruit in countries with temperate climates, including large portions of southern Brazil. Four regions account for the bulk of Brazilian apple production: Santa Catarina state, Rio Grande do Sul state, Parana state, and Sao Paulo state (Ming, 2019). In 2020, Brazil produced 983,247 tons of apples (FAO, 2022). According to the Brazilian Association of Apple Producers (ABPM), about 95% of fresh apples are sold to the Brazilian domestic market. A large portion of the processing apples, however, are exported in form of apple juice. In 2020, for example, Brazil exported 34,000 tons of apple juice (Conab, 2020).

Pests and diseases must be managed to produce high quality fresh fruit and processing apples in Brazil. Bitter rot is one of the most important diseases in Brazilian apple production areas causing losses from 9 to 15% (ABPM, 2020). It is caused by multiple *Colletotrichum* species (Munir et al., 2016; [Khodadadi et al., 2020](#); González and Sutton, 2004; Oo et al., 2018). The symptoms of apple bitter rot on fruit start with light brown, circular lesions. As the lesions expand, conidiophores rupture the fruit epidermis, forming small tufts (Henz et al., 1992). Acervuli are produced in abundance on the mature apple fruit and generate conidia that are rain splash-dispersed from one fruit to another (Peres et al., 2005; Everett et al., 2018). The conidia germinate with sufficient moisture and optimal temperature to form

apressoria. The pathogen then enters the epidermis layers to either form lesions or remains quiescent depending on the environment and maturity and apples (Wharton and Diéguez-Uribeondo, 2004; Horowitz et al., 2002; Schumann and D'Arcy, 2010).

Numerous *Colletotrichum* species were isolated from apple in Brazil, including *C. nymphaeae* and *C. melonis* of the *C. acutatum* species complex and *C. fructicola*, *C. theobromicola*, *C. alienum*, and *C. asianum* of the *C. gloeosporioides* species complex (Gelain et al., in press; Velho et al., 2014; Velho et al., 2015).

Colletotrichum is a cosmopolitan fungal genus comprised of more than 189 species distributed throughout the world. The *Colletotrichum* genus comprises nine major clades, small clusters, and isolated species, and each clade represents a species complex (Cannon et al., 2012). Among species complexes, *Colletotrichum acutatum* and *Colletotrichum gloeosporioides* are most associated with bitter rot in Brazilian apple orchards (Gelain et al., in press; Velho et al., 2014; Velho et al., 2015).

Colletotrichum acutatum species complex consists of about 30 species (Damm et al. 2012), including *Colletotrichum nymphaeae* and *Colletotrichum melonis*. They are the most common species isolated from apple fruit in Lapa, Porto Amazonas, and Parana municipalities of Parana state (Gelain et al., in press), one of the most important apple producing regions in Brazil. *C. nymphaea*, *C. fioriniae*, and *C. godetiae* from *Colletotrichum acutatum* species complex and *C. fructicola*, *C. aenigma*, *C. siamense* and *C. theobromicola* from *Colletotrichum gloeosporioides* species complex, are also known to cause bitter rot on apple in more than one country

(Jayawardena et al., 2016; Peres et al., 2005; Munir et al., 2016; Weir et al., 2012; Velho et al., 2015; Wang et al., 2015; Sato and Moriwaki, 2013).

Chemical control of bitter rot in Brazil is limited to the use of methyl benzimidazole carbamate (MBC; carbendazim, thiophanate-methyl) fungicides, captan, quinone outside inhibitor (QoI; trifloxystrobin, kresoxim-methyl, and pyraclostrobin) fungicides, and demethylation inhibitor (DMI; tebuconazole) fungicides (Agrofit, 2022). Despite these registered products, the chemical management of apple bitter rot is challenging because only a few are effective against this disease. For instance, *C. acutatum* species complex has inherent resistance to MBCs (Chung et al., 2006; Peres et al., 2004). Arguably the most effective fungicides are still the QoIs, but *Colletotrichum* spp. have a high risk of developing resistance to QoIs with many reports on resistance in *C. acutatum* species already published (Dowling et al., 2020).

Because of the limited number of effective fungicides for bitter rot control and documented QoI resistance, chemicals with other MOA, such as fluazinam and tebuconazole, may become more important regarding suppression of bitter rot. Tebuconazole is a demethylation inhibitor (DMI) fungicide that targets the sterol 14 α -demethylase (Chen et al., 2018) and has become an integral part of apple scab management in commercial orchards (Ilhan et al., 2006; Arslan, 2016), but is typically not used for apple bitter rot control. In plant and human fungal pathogens, several major mechanisms of resistance to DMIs have been described. These include: (i) target-site modification in the *CYP51* gene that result in reduced affinity of DMIs for

their target (Délye et al., 1997; Délye et al., 1998); (ii) overexpression of the *CYP51* gene (Hamamoto et al., 2000; Ma et al., 2006; Schnabel and Jones, 2001); (iii) reduced accumulation of DMIs in fungal cells through up-regulation of ATP-binding cassette (ABC) or major facilitator superfamily (MFS) transporters in the membrane (Hayashi et al. 2002; Nakaune et al. 1998; Palani and Lalithakumari 1999; Reimann and Deising 2005); and (iv) multiple *CYP51* genes generating various versions of *CYP51* proteins or reduce the fitness cost associated with the *CYP51* structure change (Ziogas and Malandrakis, 2015).

Fluazinam is an aminopyridine that inhibits glucose oxidation in a dose-dependent manner and suppresses the energy production process through direct inhibition of the adenosine triphosphate (ATP) synthetase in fungal cells (Vitoratos, 2014). The target gene or genes for fluazinam are not known but evidence suggests involvement of Mitogen-activated protein (MAP) kinase/Histidine kinase genes in resistance mechanisms. Lab mutants of *Sclerotinia sclerotiorum* with resistance to fluazinam did not reveal point mutations in the MAP/Histidine kinase *Shk-1* gene, but the gene was overexpressed in three of the nine mutants generated (Mao et al., 2018). This indicated a possible involvement of *Shk-1* in the resistance phenotype of at least some isolates. A link between fluazinam, procymidone, and fludioxonil resistance was also demonstrated in *B. cinerea* lab mutants (Shao et al., 2015), suggesting that modifications or overexpression of histidine kinase genes conferring resistance to fludioxonil may also confer resistance to fluazinam. Alternations of the MAP/Histidine kinase gene *OS-1* were shown to result in fludioxonil resistance in

Botrytis cinerea (Schnabel et al., 2021) but it is not known if this gene is involved in fluazinam resistance. To the best of our knowledge, there is no report of resistance to fluazinam in *Colletotrichum* spp.

In a previous study, *C. nymphaeae* isolates from Brazilian apple orchards were characterized for their in vitro efficacy to tebuconazole and fluazinam (Gelain et al., in press). Results indicated the presence of a range of sensitivity including isolates that were reduced-sensitive. The objective of this study was to further characterize these isolates and to investigate molecular mechanisms of reduced sensitivity.

2.2 MATERIALS AND METHODS

2.2.1 *Colletotrichum nymphaeae* isolates.

In 2017, isolates were obtained from ‘Eva’ apple fruit trees in the municipalities of Lapa, Porto Amazonas, and Palmeira, in the state of Paraná, southern Brazil (Table 1). In a previous study, *Colletotrichum* isolates had been collected from apple flowers or fruitlets with a frozen tissue-isolation method and were identified to the species level (Gelain et al., in press). A total of 11 isolates collected previously from apple fruitlets (Gelain et al., in press) with specific phenotypes were selected for this study. They included fluazinam sensitive isolates (n=3), fluazinam reduced-sensitive isolates (n=3), tebuconazole-sensitive isolates (n=3), and tebuconazole reduced-sensitive isolates (n=3). One of the twelve isolates (col139) had two phenotypes; it was reduced-sensitive to fluazinam and tebuconazole. The EC₅₀ values determined by Gelain et al. (in press) are listed in table 2.

2.2.2 In vitro sensitivity to fluazinam and tebuconazole.

An in vitro assay was performed three times to confirm the sensitivity of *Colletotrichum nymphaeae* isolates to fluazinam and tebuconazole. The single spore stocks used in this study were stored at -20°C on filter paper (Fong et al., 2000) with silica gel beads for 8 months.

The isolates were cultured on potato dextrose agar (PDA) for 8 days in the dark at 25°C (Gelain et al., in press). Mycelial disks (5 mm in diameter) from the margins of actively growing cultures were transferred onto fungicide-amended PDA in petri dishes (90 mm) with three replications for each concentration and two agar plugs per plate located 4 cm apart and equidistantly from petri dish margins. The fungicide-amended PDA had been autoclaved and cooled to 60°C before being amended either with fluazinam at 0.03, 0.1, 0.3, 1.0, and 3.0 µg /mL (Chen et al., 2016) or with tebuconazole at 0, 0.02, 0.1, 0.5, 2.5, and 12.5 µg /mL (Gang et al., 2015). The cultures were incubated at 25°C for five days in darkness before the diameter of each colony was measured in two perpendicular directions. EC₅₀ values were calculated by linear regression using R software (4.0.4, ec50estimator package). The means of EC₅₀ values of different phenotypes were compared by one-way ANOVA ($P < 0.05$).

2.2.3 Fungicide sensitivity assays using detached fruit.

Asymptomatic and organic ‘Gala’ apples of commercial maturity were purchased from a local grocery store and stored in a walk-in cooler at 4°C before

experiments were conducted. Each fruit was rinsed with tap water for 15 s and then dipped in water for 30 mins to remove any organic pesticide residues from the fruit surface. All fruit were dried in the running laminar flow hood and exposed to 5 mins of ultraviolet treatment (Janisiewicz et al., 2016). The apples were randomly placed in lidded plastic boxes (40 x 20 x 20 cm), kept in place by plastic cups, and kept at 100% relative humidity. Apples were stab-wounded with a 1 cc syringe needle to a 5 mm depth and then inoculated at the wound site with 50 μ L of a 10^5 /mL spore suspension. Spore suspensions were prepared as described previously (Suzaki et al., 2011). To verify the viability of spores, a portion of the same spore suspension (50 μ L) of each isolate was pipetted onto water agar medium (20 g/L) and spread out with a glass rod. Germination was verified 12 hrs later. Fungicide treatments were applied 12 hrs before inoculation (protective treatment) and 12 and 36 hrs after inoculation (curative treatments). Apples were sprayed with at 200 μ g/ml fluazinam, 0.5 mL / L (Secure, Syngenta Crop Protection LLC, Greensboro, NC) and tebuconazole at 67.5 μ g /mL, 0.15 g / L (Elite 45DF, Bayer CropScience, Leverkusen, Germany) following label recommended rate. Sterile water was applied to untreated inoculated control group apples. Experimental apples were kept at 25°C, 8 hrs light 16 hrs dark incubation procedure. Bitter rot lesion diameters were measured in two perpendicular directions after 7 days of incubation, and the entire experiment was performed two times.

2.2.4 DNA extraction.

Genomic DNA from *C. nymphaeae* isolates were extracted following the protocol described by Chi et al. (2009) with minor modifications. The mycelium for extraction was produced on cellophane placed on PDA plates (Cassago et al., 2002). After 7 days of 25°C in dark incubation, mycelium originating from mycelial plugs was harvested from the cellophane with autoclaved toothpicks and suspended in 500 µL of DNA extraction buffer. Fungal tissue was pulverized in centrifuge tubes using the Dremel tool with plastic tips. Samples were centrifuged for 10 minutes at 5,000 rpm and the supernatant was mixed with 300 µL of isopropanol by inverting the tubes 25 times followed by centrifugation for 10 mins at 1,2000 rpm. The supernatant was removed, and the pellet was washed with 800 µL of 70% ethanol and centrifuged for 2 mins at 1,4000 rpm. The supernatant was removed, and open-lid tubes were left in the laminar flow hood for 12 hrs (overnight) before 50 µL DNase-free water was added to suspend the DNA.

2.2.5 Primer design and PCR amplification.

CYP51A (MK990546.1), *CYP51B* (MK990546.2), and *OS-1* (XM_035483032.1) complete genome sequences and the NCBI primer blast online designing tool (National Center for Biotechnology Information, Bethesda, MD) was used for the design of primers specific to *Colletotrichum nymphaeae* gene sequences. All primer options were left on standard settings (product size range: 1000 to 1200 bp; primer size range: 0-30 nucleotides; GC content: 40-60%; and melting temperature range, 58-

61°C). For the design of *CYP51A&B* and *OS-1* genes expression analysis, settings were switched to product size: 100 to 200 bp; 0-30 nucleotides; GC content: 40-60%; and melting temperature range 60-63°C. 19 primer pair sets were designed for *CYP51A&B* and *OS-1* genes amplification, sequencing, and overexpression study (Table 2).

PCR amplifications were done in Bio-Rad Laboratories PCR Thermal Cycler T100™ (Bio-Rad Laboratories, Hercules, CA). The *CYP51A&B* genes amplification was performed with the following procedure: initial denaturation at 94°C for 10 mins, followed by 35 cycles at 94°C for 40 s, 56°C for 160s, and at 72°C for 2 mins, with final elongation at 72°C for 10 mins. The parameters were used to amplify *OS-1* gene fragment: 94°C for 40 s, 55°C for 165 s, and at 72°C for 2 mins, with final elongation at 72°C for 10 mins; 94°C for 40 s, 55°C for 2 mins, and at 72°C for 2 mins, with final elongation at 72°C for 10 mins. The PCRs were performed in a final volume of 40 µL using the AccuPower® PyroHotStart Taq PCR PreMix (Bioneer, Oakland, CA) with the following reagent: 2 µL genomic DNA (50-80 ng), 2 µL forward primer (1 µM), 2 µL reverse primer (1 µM), 36 µL nuclease-free water. After the PCR program finished, all PCR product samples were purified with Monarch® PCR & DNA Cleanup Kit (New England Biolabs, Ipswich, MA).

Table 1. Primers used for amplification of *CYP51A*, *CYP51B*, and *OS-1* gene

Primers	Sequence (5'-3')	Description
CYP51A-1	TCACAGTGATTTTCCTAGTGCTTAT	Amplify and sequence <i>CYP51A</i> gene fragment
CYP51A-8	GCCTTTCGCTTAGACCAAACCTT	
CYP51A-2	GCACGGTACTTGACGAAGAA	Sequence <i>CYP51A</i> gene fragment
CYP51A-3	AGGTTGCTAGACTTGGCTCC	Sequence <i>CYP51A</i> gene fragment
CYP51A-4	TAGGCGGTCAAGCTGTCAAT	Sequence <i>CYP51A</i> gene fragment

CYP51A-5	GTTTGGACTTACGCAAAGGCT	Sequence <i>CYP51A</i> gene fragment
CYP51A-6	TTGGCATCAACCATGCCCT	Sequence <i>CYP51A</i> gene fragment
CYP51A-7	GCAGAAGAGCTTTATCAAGAGCA G	Sequence <i>CYP51A</i> gene fragment
CYP51B-1	CAAAAGGAAGAAGAGCTTGCG	Amplify and sequence <i>CYP51B</i> gene fragment
CYP51B-8	TCTGCACTGGTTGGTGGAT	
CYP51B-2	ACTTCTTGAGGAGACCCAT	Sequence <i>CYP51B</i> gene fragment
CYP51B-3	CATTAGCTTAGCTGCTGGCTG	Sequence <i>CYP51B</i> gene fragment
CYP51B-4	CAGATTGACGACGCCAGAC	Sequence <i>CYP51B</i> gene fragment
CYP51B-5	TGCGGAGGATATCTACTGTT	Sequence <i>CYP51B</i> gene fragment
CYP51B-6	GGGCTCGGGGAAGTAGAC	Sequence <i>CYP51B</i> gene fragment
CYP51B-7	GCCAACATGATGATTGCCCT	Sequence <i>CYP51B</i> gene fragment
OS-1-1	AGCGCTGTACTTCTTACGC	Amplify <i>OS-1</i> gene fragment with primer OS-1-6 and OS-1-10, and sequence
OS-1-2	TTTTCGATTCTTTGGGCAAGC	Sequence <i>OS-1</i> gene fragment
OS-1-3	CTGCTGGTTGACCACAACCTC	Sequence <i>OS-1</i> gene fragment
OS-1-4	CTTCCAAGTACCGTCGACAC	Sequence <i>OS-1</i> gene fragment
OS-1-5	CGAGACAGGAAATCAGCAGC	Sequence <i>OS-1</i> gene fragment
OS-1-6	ACCTCTCGGGCAATCTTGG	Amplify <i>OS-1</i> gene fragment with primer OS-1-10 and sequence
OS-1-7	CTTACGATACAGGTCCGCAGT	Amplify <i>OS-1</i> gene fragment with primer OS-1-12 and sequence
OS-1-8	AGACCCAATGTCATAGACTGTTT	Sequence <i>OS-1</i> gene fragment
OS-1-9	TGTGGAATGAACTAACGGTCA	Amplify <i>OS-1</i> gene fragment with primer OS-1-16 and sequence
OS-1-10	CAGGATCGGCCTTCTTAATAGTCA	Amplify <i>OS-1</i> gene fragment with primer OS-1-1 and sequence
OS-1-11	CTGGCGAACAGTCTACTCAC	Amplify <i>OS-1</i> gene fragment with primer OS-1-16 and sequence
OS-1-12	GGTTCTTCGAGAGCAGGAATC	Amplify <i>OS-1</i> gene fragment with primer OS-1-7 and sequence
OS-1-13	CGAGACAATGCTGTCTTTGATTG	Sequence <i>OS-1</i> gene fragment
OS-1-14	GAACTATGTGACCTCAGTAGCA	Sequence <i>OS-1</i> gene fragment
OS-1-15	ATCCAGACCATTCTCAAGTGTG	Sequence <i>OS-1</i> gene fragment
OS-1-16	GAAGCTCTTAGCCACTGAAGG	Amplify <i>OS-1</i> gene fragment with primer OS-1-9 OS-1-11, and sequence
OS-1-F	AACGTCAACGCAATGGCAA	Amplify 605 bp <i>OS-1</i> gene fragment
OS-1-R	GTCGAGATACTGCGGACCTG	




Fig 1. *CYP51A* (upper schematic) and *CYP51B* (lower schematic) gene length, intron locations, and locations of primers used in this study.

2.2.6 Sequence analysis.

The sequences of *CYP51A&B* and *OS-1* genes were commercially sequenced using Eurofins USA service (Louisville, KY). Gene sequences were compared and assembled using Seqman Pro (DNASTAR Inc). Coding region nucleotide sequence alignments were done with MegAlign (DNASTAR Inc). Sequence reversions were achieved with SeqBuilder Pro (DNASTAR Inc). The positions of primers and Sanger sequence reaction directions are shown in Fig 1 and Fig 2.

2.2.7 Statistical analysis.

Statistical significance was determined in JMP Pro 15 (V15; 2019 SAS Institute Inc.) by one way ANOVA, Tukey, and least significant difference tests at $P = 0.05$. R

software version 4.0.4 with the ec50estimator R package (v0.1.0; Alves, 2020) was used to estimate the EC₅₀ values.

2.3 RESULTS

2.3.1 *Colletotrichum nymphaeae* sensitivity to fluazinam and tebuconazole.

The previously reported EC₅₀ values were similar to the EC₅₀ values obtained in this study for fluazinam and tebuconazole sensitive and reduced-sensitive isolates (Table 2; $P=0.3$). The isolates previously found to be sensitive were still sensitive and the isolates found to be categorized as reduced sensitive were still reduced sensitive. Specifically, the average ‘reduced-sensitivity’ factor for fluazinam only changed from previously 4.4 to now 4.3 and only changed for tebuconazole from previously 10.5 to now 8.8. Based on the EC₅₀ values, isolates of *C. nymphaeae* were categorized as Flu^{RS} (EC₅₀ ranging from 0.5-0.76 mg/L), Flu^S (EC₅₀ values ranging from 0.1-0.15 mg/L), Teb^{RS} (EC₅₀ values ranging from 0.22-0.69 mg/L), and Teb^S (EC₅₀ values ranging from 0.04-0.05 mg/L) (Table 1). The overall range and mean for fluazinam sensitive and reduced sensitive isolates were 0.10 to 0.64 mg/L and 0.32 mg/L, respectively (Table 2). For tebuconazole, the overall range and mean for sensitive and reduced-sensitive isolates were 0.04 to 0.69 mg/L and 0.24 mg/L, respectively (Table 2).

Table 2. Isolate names, fungicide sensitivity phenotype, geographical origin, year of isolation and EC₅₀ values obtained before and after long-term storage of

Colletotrichum nymphaeae isolates from Brazilian apple orchards

Isolate	Phenotype ^x	Origin (Municipality)	Year of isolation	Before storage ^y EC ₅₀ value (mg/L)	After storage ^z
col136	Flu ^S	Lapa	2017	0.12	0.10
col142	Flu ^S	Lapa	2017	0.15	0.13
col112	Flu ^S	Lapa	2017	0.15	0.14
col049	Flu ^{RS}	Lapa	2017	0.54	0.50
col139	Flu ^{RS}	Lapa	2017	0.56	0.41
col140	Flu ^{RS}	Porto Amazonas	2017	0.76	0.64
col134	Teb ^S	Palmeira	2017	0.04	0.04
col113	Teb ^S	Lapa	2017	0.04	0.05
col077	Teb ^S	Lapa	2017	0.04	0.05
col083	Teb ^{RS}	Palmeira	2017	0.22	0.26
col139	Teb ^{RS}	Lapa	2017	0.39	0.37
col012	Teb ^{RS}	Lapa	2017	0.66	0.69

^xFlu = fluazinam; Teb = tebuconazole; RS = Reduced Sensitive S = Sensitive.

^yDetermined by Gelain et al. in press.

^zEC₅₀ values after 8 months of storage on filter paper at -20°C.

2.3.2 Detached apple fruit assay.

The first lesions of apple bitter rot were observed within one week after inoculation and measured when approximately 24 mm wide in diameter. The Flu^S and Flu^R isolates were controlled effectively with the protective 12h treatment (Fig. 2A). However, in the curative 12h treatment the Flu^{RS} isolates were significantly harder to control compared to the Flu^S isolates. Although not statistically significant at the 0.05 level, the control efficacy was numerically lower for the Flu^{RS} isolates in the curative 36h treatment compared to the Flu^S isolate group.

Teb^S and Teb^{RS} isolates were equally well controlled when tebuconazole was applied protectively (protective 12h treatment) and curatively when tebuconazole was applied 12 h after inoculation (curative 12h treatment). However, when tebuconazole was applied 36 h after inoculation, the Teb^{RS} isolates were significantly harder to control compared to the Teb^S isolates according to the ANOVA analysis (Fig. 2B; $P \leq 0.001$).




Fig 2. Protective and curative activity of (A) fluazinam (Flu) and (B) tebuconazole (Teb) against *Colletotrichum nymphaeae* isolates with different fungicide sensitivity phenotypes on detached apple fruit. Each bar represents the mean of two independent experiments with three replicates per experiment and three isolates per phenotype. ^{RS} = Reduced sensitive isolates ^S = Sensitive isolates. Different letters indicate significant differences between treatments using Tukey's honest significant difference ($\alpha = 0.05$).

2.3.3 Sequencing of the *OS-1*, *CYP51A*, and *CYP51B* genes.

The complete *OS-1* gene contained 4 introns of 56 bp, 52 bp, 56 bp, and 65 bp length and 5 exons of 844bp, 33bp, 1094bp, 1503bp, and 441bp length (Fig. 3). The entire *OS-1* gene was 4,144 bp long and was sequenced for one sensitive (col136) and one reduced-sensitive (col139) isolate. Between the two genes there were 16 nucleotide variations in exons (15 variations). Most of these variations were silent mutations, however, there were four that led to amino acid substitutions. They were located at positions 1336, 1348, 1379, and 1516. This variable region of the gene was further amplified with primers OS-1-F and OS-1-R (Table 1) and sequenced for two more fluazinam sensitive isolates (col142 and col112) and two more fluazinam reduced-sensitive isolates (col49 and col140). Results indicated that reduced-sensitive isolates had two amino acid substitutions (L446V, A450P/S) compared to the sensitive isolates (Fig. 3).

The length of the *CYP51A* gene was 1,707 bp and contained 3 exons and 2 introns 71 bp and 75 bp in length (Fig. 1). The *CYP51B* gene length was 1,720 bp in

length and contained 2 introns of 80 bp and 65 bp in length and 3 exons (Fig. 1). No nucleotide or amino acid variations were detected in the *CYP51A* or in the *CYP51B* genes between tebuconazole sensitive isolate col134 and reduced-sensitive isolate col139.




Fig 3. *OS-1* gene length, intron locations, variable nucleotide sequence locations leading to amino acid changes, and locations of primers used in this study.

2.4 DISCUSSION

Detached fruit assays showed differential efficacy of formulated fluazinam applied at the recommended field rate against sensitive and reduced sensitive *C. nymphaeae* isolates. Like many other fungicides, fluazinam is used as a protectant and applied in the field prior to infection events (EFSA, 2019). Our study shows that in detached fruit assays protective treatments achieved high efficacy against both fluazinam phenotypes suggesting that field applications may be effective against fluazinam sensitive and reduced-sensitive phenotypes. However, our study also

showed weaknesses of curative applications, which did not control reduced-sensitive isolates as well as sensitive isolates. Curative applications of fungicides are sometimes used to prevent disease progression in the field after extensive rainfalls may have washed off previous fungicide residues on the fruit or leaf tissues (Schepers, 1996). In the field, such performance weaknesses would lead to faster relative growth of reduced-sensitive strains compared to sensitive strains and thus lead to greater speed of selection for resistant isolates (van den Bosch et al., 2011).

Detached fruit assays also show differential efficacy of tebuconazole against sensitive and reduced-sensitive *C. nymphaeae* isolates. Tebuconazole is a fungicide that can be used for both preventative and curative control of fungal diseases because it kills fungi on the surface of the plant tissue and because it is systemic, killing fungi that have already penetrated the plant epidermis. Results of this study showed that protective and 12h curative treatments had high efficacy against both sensitive and reduced-sensitive phenotypes suggesting that typical preventative applications in orchards would be effective against strains with multiple tebuconazole sensitivity phenotypes. However, our research showed weaknesses of 36h curative applications, which indicated that Teb^{RS} isolates may survive better such treatments in the field. This may lead to greater speed selection of reduced sensitive strains compared to sensitive strains.

The reduced sensitivity to fluazinam and tebuconazole was stable after 8 months of long-term storage, indicating there must be a molecular basis for the phenotypes.

Reduced sensitivity and resistance to fluazinam has been described in field isolates of

Botrytis cinerea from Japanese bean fields (Tamura et al., 2000) and in *Phytophthora infestans* from Dutch potato fields (Schepers et al., 2018), respectively. However, no information on the stability of reduced sensitivity or resistance was reported. Various studies have been performed to study DMI fungicide resistance. Reduced sensitivity to flutriafol in *Cercospora beticola* remained stable after 5 months in storage (Karaoglanidis and Thanassouloupoulos, 2002) and flusilazole resistance in *Venturia inaequalis* was found to be stable over 6 months of consecutive transfers but became unstable after 10 months (Köeller et al., 1991). Decreased resistance to flusilazole was reported for *V. inaequalis* isolates after 7 months of cold storage and in *Monilinia fructicola* isolates with reduced sensitivity to propiconazole after 34 months of cold storage (Cox et al., 2007).

Tebuconazole is registered for apple bitter rot control in Brazil, but not commonly used for this purpose; fluazinam is registered only for other diseases of apple (EPA, 2020). None of them is recommended for apple bitter rot control in the USA. The lack of effective fungicides and current resistance problems with existing fungicides (i.e. quinone outside inhibitors) may make it more difficult in the future to control bitter rot in commercial apple field production. The availability of additional options with the ability to at least suppress the disease may increase bitter rot control success and may also extend the productive life of existing products if they are not used as often.

The molecular mechanisms of resistance or reduced sensitivity to fluazinam are not yet known, but some studies suggest involvement of MAP/Histidine kinase genes.

Previous research showed the possible MAP/Histidine kinase *Shk-1* gene, involved in osmotic signal transduction, involvement in fluazinam resistance (Mao et al., 2018). Specifically, the authors found the *Shk-1* gene overexpressed in some but not all *Sclerotinia sclerotiorum* isolates from oilseed rape resistant to fluazinam (Mao et al., 2018). The sequencing of the *Shk-1* homologue of *C. nymphaeae* reported in this study provides the molecular basis for further examination of expression patterns in isolates with reduced sensitivity to fluazinam. Although point mutations in MAP/Histidine Kinase genes have not been found to confer resistance to fluazinam, our study provides preliminary evidence in form of three amino acid variations between sensitive and reduced sensitive isolates. More studies are warranted to confirm the relationship between these amino acid alterations and the reduced-sensitivity phenotype.

In summary, *C. nymphaeae* from apples appear to have natural resistance (reduced sensitivity) to fluazinam and tebuconazole. Fluazinam and tebuconazole inhibited mycelial growth of *C. nymphaeae* in vitro and was as effective in reducing apple bitter rot development on detached fruits. The apple growers will benefit from fluazinam and tebuconazole application and registration. With limited number of fungicides available for apple bitter rot management and the risk of fungicide resistance selection, further studies with other alternative fungicides are still necessary.

Literature Cited

- Abbott, C. P., and Beckerman, J. L. 2018. Incorporating adjuvants with captan to manage common apple diseases. *Plant Dis.* 102:231-236.
- Abid, M., Chohan, S., Perveen, R., and Anees, M. 2019. First report of bitter rot of apple caused by *Colletotrichum siamense* in Pakistan. *Plant Dis.* 3:583.
- ABPM. Associação Brasileira de Produtores de Maçã. 2020. Available at: <https://www.abpm.org.br/>. [Accessed May 5, 2022].
- Agrofit. 2022. Sistema de Agrotóxicos Fitossanitários. Available at: http://agrofit.agricultura.gov.br/agrofit_cons/principal_agrofit_cons. [accessed 2022 May 5].
- Afanador-Kafuri, L., M. Maymon, D. M., and Freeman, S. 2003. Characterization of *Colletotrichum* isolates from tamarillo, passiflora and mango in Colombia and identification of a unique species from the genus. *Phytopathology* 93:579–587.
- Alves, K. S. 2020. ec50estimator: An automated way to estimate EC₅₀ for stratified datasets. R package version 0.1.0. <https://CRAN.R-project.org/package=ec50estimator>.
- Arslan, U. 2016. Efficacy of boric acid, monopotassium phosphate and sodium metabisulfite on the control of apple scab. *J. Phytopathol.* 164:678-685.
- Avila-Adame, C., Olaya, G., and Köller, W. 2003. Characterization of *Colletotrichum graminicola* isolates resistant to strobilurin-related QoI fungicides. *Plant Dis.* 87:1426-1432.

- Beever, R. E., Olsen, T. L., and Parkes, S. L. 1995. Vegetative compatibility groups in *Colletotrichum gloeosporioides* (*Glomerella cingulata*) from apple and other fruits. *Aust. J. Plant Pathol.* 24:126–132.
- Bernstein, B., Zehr, E. I., and Dean, R. A. 1995. Characteristics of *Colletotrichum* from peach, apple, pecan, and other hosts. *Plant Dis.* 79:478-482.
- Bogo, A., Casa, R. T., Agostineto, L., Gonçalves, M. J., and Rufato, L. 2012. Effect of hail protection nets on apple scab in ‘Royal Gala’ and ‘Fuji’ apple cultivars. *Crop Prot.* 38:49-52.
- Børve, J., and Stensvand, A. 2007. *Colletotrichum acutatum* found on apple buds in Norway. *Plant Health Prog.* 8:49.
- Cannon, P. F., Damm, U., Johnston, P. R., and Weir, B. S. 2012. *Colletotrichum*: current status and future directions. *Stud. in Mycol.* 1:181-213.
- Carraro, T. A., Moreira, R. R. Gelain, J., May-De Mio, L. L. 2022. Etiology and epidemiology of diseases caused by *Colletotrichum* spp. in persimmon, apple, peach, and grapevine. *A Revisão Anual de Patologia de Plantas.* 28:136-162.
- Cassago, A., Panepucci, R. A., Baião, and A. M. T., Henrique-Silva, F. 2002. Cellophane based mini-prep method for DNA extraction from the filamentous fungus *Trichoderma reesei*. *BMC Microbiol.* 18:14.
- Cechi, A., Ivey, M. L. L., Kaufman, R. R., Bryson, K. P., and Schnabel, G. 2020. Quinone outside inhibitor-resistant *Colletotrichum nymphaeae* isolates from strawberry lack mutations in *cytb* gene. *J. Plant Pathol.* 102:681–683.

- Chen, S., Wang, Y., Schnabel, G., Peng, C. A., Lagishetty, S., Smith, K., Luo, C., and Yuan, H. 2018. Inherent resistance to 14 α -Demethylation inhibitor fungicides in *Colletotrichum truncatum* is likely linked to *CYP51A* and/or *CYP51B* Gene Variants. *Phytopathology* 108:1263-1275.
- Chen, S. N., Luo, C. X., Hu, M. J., and Schnabel, G. 2016. Sensitivity of *Colletotrichum* species, including *C. fioriniae* and *C. nymphaeae*, from peach to demethylation inhibitor fungicides. *Plant Dis.* 100: 2434–2441.
- Chung, W., Ishii, H., Nishimura, K., Fukaya, M., Yano, K., and Kajitani, Y. 2006. Fungicide sensitivity and phylogenetic relationship of anthracnose fungi isolated from various fruit crops in Japan. *Plant Dis.* 90:506-512.
- Chi, M. H., Park, S. Y. and Lee, Y. H. 2009. A quick and safe method for fungal DNA extraction. *The Plant Pathol.* 25:108-111.
- Conab. Companhia Nacional de Abastecimento. 2020. Available at: <https://www.conab.gov.br/info-agro/safras/graos/boletim-da-safra-de-graos/>. [Accessed May 5, 2022].
- Cox, K. D., Bryson, P. K., and Schnabel, G. 2007. Instability of propiconazole resistance and fitness in *Monilinia fructicola*. *Phytopathology* 97:448-53.
- Crouch, J., O'Connell, R., Gan, P., Buiate, E., Torres, M. F., Beirn, L., Shirasu, K., and Vaillancourt, L. 2014. Genomics of Plant-Associated Fungi: Monocot Pathogens. *The Genomics of Colletotrichum*, eds. Dean, R., Lichens-Park, A., Kole, C. Cham: Springer, Berlin, Heidelberg, p.69-102.

- Daykin, M. E. and Milholland, R. D. 1984. Ripe rot of muscadine grape caused by *Colletotrichum gloeosporioides* and its control. *Phytopathology* 74:710.
- Damm, U., Cannon, P. F., Woudenberg, J. H., and Crous, P. W. 2012. The *Colletotrichum acutatum* species complex. *Stud Mycol.* 73:37-113.
- Dean, R., Kan, Jan a. L. Van, Pretorius, Z. A., Hammond-Kosack, K. E., Pietro, A. D., Spanu, P. D., Rudd, J. J., Dickman, M., Kahmann, R., Ellis, J., and Foster, G. D. 2012. The Top 10 fungal pathogens in molecular plant pathology. *Mol. Plant Pathol.* 13:414-430.
- Denardi, F., Berton, O., and Spengler, M. M. 2003. Resistência genética à podridão amarga em maçãs, determinada pela taxa de desenvolvimento da doença em frutos com e sem ferimentos. *Revista Brasileira de Fruticultura.*25:494-497.
- Délye, C., Laigret, F., and Corio-Costet, M. F. 1997. Cloning and sequence analysis of the eburicol 14 α -demethylase gene of the obligate biotrophic grape powdery mildew fungus. *Gene.*195:29-33.
- Délye, C., Bousset, L., and Corio-Costet, M. F. 1998. PCR cloning and detection of point mutations in the eburicol 14 α -demethylase (*CYP51*) gene from *Erysiphe graminis* f. sp. *hordei*, a "recalcitrant" fungus. *Curr. Genet.* 34:399-403.
- Dowling, M., Peres, N., Villani, S., and Schnabel G. 2020. Managing *Colletotrichum* on fruit crops: A "complex" challenge. *Plant Dis.* 104: 2301-2316.
- Dzhangaliev, A. D. 2002. The Wild Apple Tree of Kazakhstan. Pages 63-303 in: *Horticultural Reviews.* John Wiley & Sons, Ltd.

Eastburn, D. M. and Gubler, W. D. 1990. Strawberry anthracnose: detection and survival of *Colletotrichum acutatum* in soil. *Plant Dis.* 74: 161–163.

EFSA. 2019. Public consultation on the active substance Fluazinam. Available at: <https://www.efsa.europa.eu/en/consultations/call/public-consultation-active-substance-fluazinam> [Accessed Aug 5, 2022].

EPA. 2020. Pesticide Product Label, AX FLUAZINAM. Available at: https://www3.epa.gov/pesticides/chem_search/ppls/089167-00074-20200626.pdf [Accessed Aug 5, 2022].

Everett, K. R., Pushparajah, I. P. S., Timudo, O. E., Ah Chee, A., Scheper, R. W. A., Shaw, P. W., Spiers, T. M., Taylor, J. T., Wallis, D. R., and Wood, P. N. 2018. Infection criteria, inoculum sources and splash dispersal pattern of *Colletotrichum acutatum* causing bitter rot of apple in New Zealand. *Eur. J. Plant Pathol.* 152: 367–383.

FAO. Food and Agriculture Organization of United Nations. 2022. Crops and livestock products. <https://www.fao.org/faostat/en/#data/QCL> [Accessed May 5, 2022].

Fong, Y. K., Anuar, S., Lim, H. P., Tham, F. Y., and Sanderson, F. R. 2000. A modified filter paper technique for long-term preservation of some fungal cultures. *Mycologist* 14:127–130.

- Forcelini, B. B. and Peres, N. A. 2018. Widespread resistance to QoI fungicides of *Colletotrichum acutatum* from strawberry nurseries and production fields. *Plant Health Prog.* 19:338-341.
- Freeman, S. 2008. Management, Survival Strategies, and Host Range of *Colletotrichum acutatum* on Strawberry. *HortScience.* 43:66-68.
- Freeman, S., Shalev, Z., and Katan, J. 2002. Survival in soil of *Colletotrichum acutatum* and *C. gloeosporioides* pathogenic on strawberry. *Plant Dis.* 86:965–970.
- Forsline, P. L., Aldwinckle, H. S., Dickson, E. E., Luby, J. J. and Hokanson, S.C., 2003. Collection, maintenance, characterization, and utilization of wild apples of Central Asia. *Horticultural Reviews-Westport Then New York*, p.1-62.
- Gang, G. H., Cho, H. J., Kim, H. S., Kwack, Y. B., and Kwak, Y. S. 2015. Analysis of fungicide sensitivity and genetic diversity among *Colletotrichum* species in sweet persimmon. *The Plant Pathol. J.* 31:115–122.
- Gelain, J., Lykins, S., Rosa, P. F., Soares, A. T., Dowling, W., Schnabel, G., May De Mio, L. L. 2022. Identification and fungicide sensitivity of *Colletotrichum* spp. from apple flowers and fruitlets in Brazil. *Plant Dis.* in press.
- González, E. and Sutton, T. B. 2004. Population diversity within isolates of *Colletotrichum* spp. causing Glomerella leaf spot and bitter rot of apples in three orchards in North Carolina. *Plant Dis.* 88:1335-1340.
- Guo, Z., Miyoshi, H., Komyoji, T., Haga, T., and Fujita, T. 1991. Uncoupling activity of a newly developed fungicide, fluazinam [3-chloro-N-(3-chloro-2,6-dinitro-4-

trifluoromethylphenyl)-5-trifluoromethyl-2-pyridinamine]. *Bioenergetics*. 1056:89-92.

Hamada, N. A., Moreira, R. R., Figueiredo, J. A. G., and Mio, L. L. M. De. 2020.

Colletotrichum acutatum complex isolated from apple flowers can cause bitter rot and Glomerella leaf spot. *Bragantia*. 79:399-406.

Hamamoto, H., Hasegawa, K., Nakaune, R., Young, J. L., Makizumi, Y., Akutsu, K., and Hibi, T. 2000. Tandem repeat of a transcriptional enhancer upstream of the sterol 14 α -demethylase gene (*CYP51*) in *Penicillium digitatum*. *Appl. Environ. Microbiol.* 66:3421-3426.

Hayashi, K., Schoonbeek, H. J., and De Waard, M. A. 2002. Expression of the ABC transporter *BcatrD* from *Botrytis cinerea* reduces sensitivity to sterol demethylation inhibitor fungicides. *Pestic. Biochem. Physiol.* 73:110-121.

Henz, G. P., Boiteux, L. S., and Lopes, C. A. 1992. Outbreak of strawberry anthracnose caused by *Colletotrichum acutatum* in central Brazil. *Plant Dis.* 76:212.

Horowitz, S., Freeman, S., and Sharon, A. 2002. Use of green fluorescent protein-transgenic strains to study pathogenic and nonpathogenic lifestyles in *Colletotrichum acutatum*. *Phytopathology* 92:743-749.

Howard, C. M. 1972. A strawberry fruit rot caused by *Colletotrichum fragariae*. *Phytopathology* 62:600-602.

- Hu, M., Grabke, A., Dowling, M. E., Holstein, H. J., and Schnabel, G. 2015. Resistance in *Colletotrichum siamense* from peach and blueberry to thiophanate-methyl and azoxystrobin. *Plant Dis.* 99:806-814.
- Hyde, K. D., Cai, L., Mckenzie, E. H. C., Yang, Y. L., Zhang, J. Z., and Prihastuti, H. 2009. *Colletotrichum*: A catalogue of confusion. *Fungal Divers.* 39:1-17.
- Ivey, M. L. L. 2016. Bitter rot of apple. Available at: <https://ohioline.osu.edu/factsheet/plpath-fru-20> [Accessed July 27, 2022].
- Ilhan, K., Arslan, U., and Karabulut, O. A. 2006. The effect of sodium bicarbonate alone or in combination with a reduced dose of tebuconazole on the control of apple scab. *Crop Protect.* 25:963-967.
- Janisiewicz, W. J., Takeda, F., Nichols, B., Glenn, D. M., II, W. M. J., and Camp, M. J. 2016. Use of low-dose UV-C irradiation to control powdery mildew caused by *Podosphaera aphanis* on strawberry plants. *Can. J. Plant Pathol.* 38:430-439.
- Jayawardena, R. S., Hyde, K. D., Damm, U, Cai, L., Liu, M., Li, X. H., Zhang, W., Zhao, W. S., and Yan, J. Y. 2016. Notes on currently accepted species of *Colletotrichum*. *Mycosphere.* 7:1192–1260.
- Jones, A. L., Ehret, G. R., Meyer, M. P., and Shane, W. W. 1996. Occurrence of bitter rot on apple in Michigan. *Plant Dis.* 80:1294-1297.
- Karaoglanidis, G. S. and Thanassoulopoulos, C. C. 2002. Phenotypic Instability of *Cercospora beticola* Sacc. Strains expressing resistance to the sterol demethylation-

inhibiting (DMI) fungicide flutriafol after cold exposure. *J. Phytopathology* 150: 692-696.

Katsurayama, Y., Boneti, J. I. S., and Becker, W. F. 2000. Mancha foliar da Gala: Principal doença de verão da cultura da macieira. *Agropecuária* 13:14-19.

Khodadadi, F., González, J. B., Martin, P. L., Giroux, E., Bilodeau, G. J., Peter, K. A., Doyle, V. P., and Aćimović, S. G. 2020. Identification and characterization of *Colletotrichum* species causing apple bitter rot in New York and description of *C. noveboracense* sp. nov. *Sci. Rep.* 1:1-19.

Kim, Y., Dixon, E. W., Vincelli, P., and Farman, M. L. 2003. Field resistance to Strobilurin (QoI) fungicides in *Pyricularia grisea* caused by mutations in the mitochondrial *Cytochrome b* Gene. *Phytopathology*. 93:891-900.

Köller, W., Smith, F. D. and Reynolds, K. L. 1991. Phenotypic instability of flusilazole sensitivity in *Venturia inaequalis*. *Plant Pathol.* 40: 608-611.

Lesniak, K. E., Proffer, T. J., Beckerman, J. L., and Sundin, G. W. 2011. Occurrence of QoI Resistance and Detection of the G143A Mutation in Michigan Populations of *Venturia inaequalis*. *Plant Dis.* 95:927-934.

Ma, Z., Proffer, T. J., Jacobs, J. L., and Sundin, G. W. 2006. Overexpression of the 14 α -demethylase gene (*CYP51*) mediates fungicide resistance in *Blumeriella jaapii*. *Appl. Environ. Microbiol.* 72:2581-2585.

- MacKenzie, S. J. and Peres, N. A. 2012. Use of leaf wetness and temperature to time fungicide applications to control anthracnose fruit rot of strawberry in Florida. *Plant Dis.* 96:522-528.
- Martin, P. L., Krawczyk, T., Pierce, K., Thomas, C., Khodadadi, F., Aćimović, S. G., and Peter, K. A. 2022. Fungicide sensitivity of *Colletotrichum* species causing bitter rot of apple in the mid-Atlantic U.S.A. *Plant Dis.* 106: 549-563.
- Moral, J., Oliveira, R., and Trapero, A. 2009. Elucidation of the disease cycle of olive anthracnose caused by *Colletotrichum acutatum*. *Phytopathology* 99:548-556.
- Moreira, R. R., Zielinski, E. C., Castellar, C., Bergamin Filho, A., and May-De Mio, L. L. 2020. Study of infection process of five species of *Colletotrichum* comparing symptoms of Glomerella leaf spot and bitter rot in two apple cultivars. *Eur. J. Plant Pathol.* 159:37-53.
- Mao, X. W., Li, J. S., Chen, Y. L., Song, X.S., Duan, Y. B., Wang, J. X., Chen, C. J., Zhou, M. G., Hou, Y. P. 2018. Resistance risk assessment for fluazinam in *Sclerotinia sclerotiorum*. *Pestic. Biochem. and Phys.* 144:27-35.
- Munir, M., Amsden, B., Dixon, E., Vaillancourt, L., and Gauthier, N. A. W. 2016. Characterization of *Colletotrichum* species causing bitter rot of apple in Kentucky orchards. *Plant Dis.* 100:2194–2203.
- Nakaune, R., Adachi, K., Nawata, O., Tomiyama, M., Akutsu, K. and Hibi, T. 1998. A novel ATP-binding cassette transporter involved in multidrug resistance in the

phytopathogenic fungus *Penicillium digitatum*. Appl. Environ. Microbiol. 64:3983–3988.

Nekoduka, S., Tanaka, K., and Sano, T. 2018. Epidemiology of apple bitter rot caused by *Colletotrichum acutatum sensu lato*. J. Gen. Plant Pathol. 84:262-271.

Oh, J. and Kang, N. 2002. Efficacy of fluazinam and iprodione+propineb in the suppression of *Diaporthe phaseolorum*, *Colletotrichum truncatum* and *Cercospora kikuchii*, the causal agents of seed decay in soybean. The Plant Pathol. J. 18:216-220.

Oliveira, M. S., Cordova, L. G., and Peres, N. A. 2020. Efficacy and baseline sensitivity of succinate-dehydrogenase-inhibitor fungicides for management of *Colletotrichum* crown rot of strawberry. Plant Dis. 104:2860-2865.

Oo, M. M., Yoon, H., Jang, H. A., and Oh, S. 2018. Identification and characterization of *Colletotrichum* species associated with bitter rot disease of apple in South Korea. Plant Pathol. J. 34:480-489.

Palani, P. V. and Lalithakumari, D. 1999. Resistance of *Venturia inaequalis* to the sterol biosynthesis-inhibiting fungicide, penconazole [1-(2-(2,4-dichlorophenyl)pentyl)-1H-1,2,4-triazole]. Mycol. Res. 103:1157–1164.

Peres, N. A., Timmer, L. W., Adaskaveg, J. E., and Correll, J. C. 2005. Lifestyles of *Colletotrichum acutatum*. Plant Dis. 89:784–796.

Peres, N. A., Souza, N. L., Peever, T. L., and Timmer, L. W. 2004. Benomyl sensitivity of isolates of *Colletotrichum acutatum* and *C. gloeosporioides* from citrus. Plant Dis. 88:125-130.

- Poti, T., Mahawan, K., Cheewangkoon, R., Arunothayanan, H., Akimitsu, K., and Nalumpang, S. 2020. Detection and molecular characterization of carbendazim-resistant *Colletotrichum truncatum* isolates causing anthracnose of soybean in Thailand. *J. Phytopathol.* 168:267-278.
- Reimann, S. and Deising, H. B. 2005. Inhibition of efflux transporter-mediated fungicide resistance in *Pyrenophora tritici-repentis* by a derivative of 4'-hydroxyflavone and enhancement of fungicide activity. *Appl. Environ. Microbiol.* 71: 3269–3275.
- Roberts, J. W. 1918. The Sources of Apple Bitter-rot Infections. U.S. Department of Agriculture, Washington, D.C.
- Sato, T. and Moriwaki, J. 2013. Re-classification of the *Colletotrichum acutatum* group A2 and A4 and re-identification of strains belonging to the groups in Japan. *Plant Protec.* 67:367–372.
- Schepers, H. T. A. M. 1996. Effect of rain on efficacy of fungicide deposits on potato against *Phytophthora infestans*. *Potato Res.* 39: 541–550.
- Schepers, H. T. A. M., Kessel, G. J. T., Lucca, F. M., Förch, G., van den Bosch, G. B. M., Topper, C. G., and Evenhuis, A. 2018. Reduced efficacy of fluazinam against *Phytophthora infestans* in the Netherlands. *Eur. J. Plant Pathol.* 151:947–960.
- Schnabel, G., and Jones, A. L. 2001. The 14 α -Demethylase (*CYP51A1*) gene is overexpressed in *Venturia inaequalis* strains resistant to myclobutanil. *Phytopathology* 91:102-110.

- Schnabel, G., Tan, Q., Schneider, V., and Ishii, H. 2021. Inherent tolerance of *Colletotrichum gloeosporioides* to fludioxonil. *Pesticide Biochem. and Phys.* 172:104767.
- Schumann, G. L. and D'Arcy, C. J. 2010. *Essential Plant Pathology*. American Phytopathological Socie. pp. 197.
- Shao, W., Ren, W., Zhang, Y., Hou, Y., Duan, Y., Wang, J., Zhou, M., and Chen C., 2015. Baseline sensitivity of natural populations and characterization of resistant strains of *Botrytis cinerea* to fluazinam. *Australas. Plant Pathol.* 44:375–383.
- Silva-Junior, G. J., Spósito, M. B., Marin, D. R., and Amorim, L. 2014. Efficacy and timing of application of fungicides for control of citrus postbloom fruit drop. *Crop Protect.* 59:51-56.
- Sinclair, J. B. 1991. Latent infection of soybean plants and seeds by fungi. *Plant Dis.* 75:220-224.
- Stensvand, A., Børve, J., and Talgø, V. 2017. Overwintering diseased plant parts and newly infected flowers and fruit as sources of inoculum for *Colletotrichum acutatum* in sour cherry. *Plant Dis.* 101:1207-1213.
- Suzaki, K. 2011. Improved method to induce sporulation of *Colletotrichum gloeosporioides*, causal fungus of grape ripe rot. *J. Gen. Plant Pathol.* 77:81–84.
- Tamura, O. 2000. Resistance development of gray mold on beans towards fluazinam and relevant countermeasures. *Abstr. 10th Symp. Res. Comm. Fungic. Resist.* Okoyama, Japan. p. 7-16

- Taylor, J. 1971. Epidemiology and symptomatology of apple bitter rot. *Phytopathology*. 61:1028-1029
- Timmer, L. W., Brown, G. E., and Zitko, S. E. 1998. The role of *Colletotrichum* spp. in postharvest anthracnose of citrus and survival of *C. acutatum* on fruit. *Plant Dis.* 82:415-418.
- Ming, P. 2019. Annual Fresh Deciduous Fruit Report. Available at: https://usdabrazil.org.br/wp-content/uploads/2021/09/Annual-Fresh-Deciduous-Fruit-Report_Brasilia_Brazil_12-19-2019.pdf [Accessed May 5, 2022].
- van den Bosch, F., Paveley, N., Shaw, M., Hobbelen, P. and Oliver, R. 2011. The dose rate debate: does the risk of fungicide resistance increase or decrease with dose?. *Plant Pathol.* 60:597-606.
- Velho, A. C., Stadnik, M. J., Casanova, L., Mondino, P., and Alaniz, S. 2014. First report of *Colletotrichum nymphaeae* causing apple bitter rot in southern Brazil. *Plant Dis.* 98:567-567.
- Velho, A. C., Alaniz, S., Casanova, L., Mondino, P., and Stadnik, M. J. 2015. New insights into the characterization of *Colletotrichum* species associated with apple diseases in southern Brazil and Uruguay. *Fungal Biol.* 4:229-244.
- Vitoratos, A. G. 2014. Mode of action and genetic analysis of resistance to fluazinam in *Ustilago maydis*. *J. Phytopathol.* 162:737-746.
- Wharton, P. S. and Diéguez-Uribeondo, J. 2004. The biology of *Colletotrichum acutatum*. *An. Jard. Bot. Madr.* 61:3-22.

Wang, B., Li, B. H., Dong, X. L., Wang, C. X., and Zhang, Z. F. 2015. Effects of temperature, wetness duration, and moisture on the conidial germination, infection, and disease incubation period of *Glomerella cingulata*. *Plant Dis.* 99:249-256.

Weir, B. S., Johnston, P. R., and Damm, U. 2012. The *Colletotrichum gloeosporioides* species complex. *Stud. Mycol.* 73:115–180.

Yokosawa, S., Eguchi, N., Kondo, K., and Sato, T. 2017. Phylogenetic relationship and fungicide sensitivity of members of the *Colletotrichum gloeosporioides* species complex from apple. *J. Gen. Plant Pathol.* 83: 291–298.

Ziogas, B. N. and Malandrakis, A. A. 2015. Sterol biosynthesis inhibitors: C14 demethylation (DMIs). p. 199-216 in: *Fungicide Resistance in Plant Pathogens*. Springer, Tokyo.